


INITIAL CORE DESCRIPTIONS

DEEP SEA DRILLING PROJECT

LEG 41

EASTERN NORTH ATLANTIC



Prepared for the
NATIONAL SCIENCE FOUNDATION
National Ocean Sediment Coring Program
Under Contract C-482
By the
UNIVERSITY OF CALIFORNIA
Scripps Institution of Oceanography
Prime Contractor for the Project

MGG 15055042

UNIVERSITY OF CALIFORNIA, SAN DIEGO

BERKELEY • DAVIS • IRVINE • LOS ANGELES • RIVERSIDE • SAN DIEGO • SAN FRANCISCO

SANTA BARBARA • SANTA CRUZ



SCRIPPS INSTITUTION OF OCEANOGRAPHY

POST OFFICE BOX 1529
LA JOLLA, CALIFORNIA 92037

Dear Colleague:

This document has been printed and distributed by the Deep Sea Drilling Project for the purpose of sample selection by interested earth scientists, sample requests being honored one year after completion of the cruise on which the samples were collected. It is an interim and informal document consisting of site data and sedimentologic and paleontologic data as known six (6) months post-cruise. These data, while completely adequate for almost all sample selection needs, will be subject to possible slight change by the time of issue of the formal cruise report, the corresponding volume of the Initial Reports of the Deep Sea Drilling Project.

The information contained herein is preliminary and privileged, consequently this document is not to be cited or used as the basis of other publications. Data cited or used in a manuscript will be considered a breach of professional ethics.

Thank you for your interest in the Deep Sea Drilling Project.

Sincerely,

N. Terence Edgar
N. Terence Edgar
Chief Scientist
Deep Sea Drilling Project

NTE:eb

INITIAL CORE DESCRIPTION (ICD)

DEEP SEA DRILLING PROJECT

LEG 41

February 17, 1975 — April 10, 1975

A Project Planned by and Carried Out With the Advice of the
JOINT OCEANOGRAPHIC INSTITUTIONS FOR DEEP EARTH SAMPLING (JOIDES)

MEMBER ORGANIZATIONS

Lamont-Doherty Geological Observatory, Columbia University
Rosenstiel School of Marine and Atmospheric Science, University of Miami
Scripps Institution of Oceanography, University of California
University of Washington
Woods Hole Oceanographic Institution
P. P. Shirshov Institute of Oceanology, Moscow, USSR
Bundesanstalt für Bodenforschung, Hannover, FRG

PARTICIPATING SCIENTISTS

Yves Lancelot, Eugen Seibold, Pavel Cepek,
Walter E. Dean, Vladislav Eremeev, James V. Gardner,
Lubomir F. Jansa, David Johnson, Valeri Krashneninnikov,
Uwe Pflaumann, J. Graham Rankin, Peter Trabant

INITIAL CORE DESCRIPTION - LEG 41

INTRODUCTION

GLOMAR CHALLENGER left Abidjan, Ivory Coast on February 17, 1975, and docked at Malaga, Spain on April 10, after having drilled five sites. The main objectives of this cruise were to obtain as complete as possible a record of the sedimentary history of the eastern North Atlantic and its margin in order to understand their evolution and compare that to the evolution of the western Atlantic and the South Atlantic.

During the planning phase for this leg, a deliberate choice was made to drill only a small number of holes, but carefully select them so that their results would yield a comprehensive picture of the different environments represented in the area. Restricting the number of holes appeared to be the best way to get deep penetration and to obtain a complete record of all the sediment facies present. In order to achieve these goals, eight primary sites, located in the deep basins, on different types of rises, and on the continental slope, were selected by the JOIDES Atlantic Advisory Panel. All the sites were selected in areas where abundant seismic reflection profiles are available thus insuring that each site could be located in a region truly representative of the general stratigraphy of the eastern North Atlantic. This also allows the drilling results to be utilized for basin-wide correlations. Much of the data obtained during Leg 41 should eventually be of more than regional interest; they bear on fundamental problems related to the

evolution of the Atlantic and more generally of a "rifted" ocean and its margins.

Of the five sites drilled during this Leg (Figure 1), two were deep basin sites (367 and 370), two were rise sites (366 and 368) and one was a continental slope site (369). The sites drilled in the basins are located on old oceanic crust, within the magnetic quiet zone, so that only the oldest part of the record was expected to provide a good stratigraphic control. The younger sediments should have been mainly deposited below the CCD.

The younger record was investigated by drilling on the rise sites. Sierra Leone Rise is a typical aseismic oceanic rise, covered with a blanket of pelagic sediments. Cape Verde Rise was also expected to be covered with pelagic sediments; however, the occurrence of well-stratified parallel reflectors on the seismic reflection profiles suggests that hemipelagic sedimentation under the influence of bottom currents could have played an important role in the building of the rise. In that respect, the site on Cape Verde Rise (363) could be considered as intermediate between a basin site and the site on the Sierra Leone Rise (366). The site on the continental slope (369), south of the Atlas region which was folded in the Tertiary, can also provide a good comparison with the sites drilled on the rises.

In addition to covering a wide range of water depths, the five Leg 41 sites also cover a wide range of latitudes, from about 5°N to about 33°N, providing control on both vertical and latitudinal environment gradients reflecting both the biogenic and terrigenous contribution to the sediments.

6

A brief description of the preliminary scientific results appeared
in Geotimes, July 1975, v. 20, (p. 18-21).




Figure 1. Leg 41 Drill Sites.

EXPLANATORY NOTES

The purpose of this volume is to present descriptions of all the cores recovered on DSDP Leg 41 to aid potential investigators in the selection of samples. The descriptions of lithology and biostratigraphy should be regarded as tentative because shore-based investigations were just beginning at the time this volume was published. The following material is intended as an aid in understanding the terminology, labeling, and the numbering conventions used on DSDP Leg 41 material.

Numbering and Depth Conventions

A site number refers to a single hole or group of holes drilled in essentially the same position using the same acoustic beacon. The first hole at a site is given the number of the site. Second or subsequent holes drilled after withdrawing from the first hole and redrilling were labeled "A", "B", etc. (e.g. Site 366A).

A core is taken by dropping a core barrel down the drill string and coring for 9.5 meters as measured by lowering of the drill string. The sediment is retained in a plastic liner 9.28 meters long inside the core barrel and in a 0.20 meter long core catcher assembly below the liner. The liner is not usually full when retrieved.

Upon recovery the liner is cut into sections of 1.5 meters, measured from the lowest point of sediment within the liner. In general the top of the core does not coincide with the top of a

section. The sections are labeled from 1 for the top (often incomplete) section to a figure as high as 6 for the bottom (complete) section, depending on the total length of core recovered. By convention, when partial recovery results, the recovered sediment is assumed to represent the top of the cored sequence. The core catcher represents sediment immediately below the lowest section.

An example of accepted convention for a sample number is "41-369-3-1 (10-20 cm)." This sample represents sediment from Leg 41, Site 369, Core 3, in Section 1 between 10 and 20 centimeters.

Handling of Cores

After a core section was cut, sealed, and labeled, it was brought into the core laboratory for processing. Sonic velocity measurements, using a Hamilton frame velocimeter, and physical property data were taken on selected samples. Samples were taken from suitable cores for inorganic and organic geochemical analysis. One of the split halves was designated a working half. Samples, including those for grain size, X-ray mineralogy, water content, carbon-carbonate, and samples for shipboard and shorebased studies of nannoplankton, foraminifera, radiolarians, diatoms, and silico-flagellates or other paleontological studies were taken.

The other half of the core section was designated an archive half. The color, texture, structure, and composition of the various lithologic units within a section were described on standard visual core description sheets (one per section) and any remarkable features

noted. Smear slides were made for each sediment of distinct lithology, microscopic examinations were made, and the descriptions were recorded. The archive half of the core section was then photographed. After the above process, both halves were stored in the ship's cold storage.

All samples now reside in cold storage at the DSDP East Coast Repository at Lamont-Doherty Geological Observatory, Palisades, New York and are available to investigators.

Sediment Analyses

Carbon-Carbonate

Organic carbon and calcium carbonate analyses were performed using a Leco 70-Second Analyzer following procedures outlined in Volumes 9 and 18 of the Initial Reports of the Deep Sea Drilling Project. Accuracy and precision of the results are as follows:

Total carbonate	$\pm 0.3\%$ (absolute)
Organic carbon	$\pm 0.06\%$ (absolute)
CaCO_3	$\pm 3\%$ (absolute)

Analyses designated "CaCO₃ bomb" were made using a method of measuring the CO₂ pressure after acidification of the sample, described by Müller and Gastner (1971). The state the accuracy of this method is $\pm 1\%$ CaCO₃.

The size classification used here is that of Shepard (1954) with the sand, silt, and clay boundaries based on the Wentworth (1922) scale: sand from 2000 to 62.5 microns, silt 62.5 to 3.91 microns, and clay less than 3.91 microns.

Standard sieve and pipette methods were used to determine the grain size distribution. The sand-size fraction was removed by wet sieving using a 63-micron sieve, and the silt and clay fractions were analyzed by standard pipette analysis. Sampling depths and volumes were calculated using equations derived from the Stokes settling velocity equation (Krumbein and Pettijohn, 1938, 95-96).

Sediment Classification

The sediment classification used on Leg 41 is outlined below. The sediment was examined on a smear slide and placed in one of the categories.

CLASSIFICATION AND NOMENCLATURE RULES

- I. Rules for class limits and sequential listing of constituents in a sediment name.
 - A. Major constituents
 1. Sediment assumes names of those constituents present in major amounts (major defined as >25%). See example in rule 1A3.
 2. Where more than one major constituent is present, the one in greatest abundance is listed farthest to the right. The remaining major constituents are listed progressively farther to the left in order of decreasing abundance.
 3. When two or more major constituents are present, class limits are based on percentage intervals:
0-5, 5-25, 25-75, 75-100.

11

B. Minor constituents

Constituents present in the amounts of 5-25% are prefixed on the sediment name by the term -bearing.

Example: 50% nannofossils, 30% radiolarians, 20% zeolites would be called a zeolite-bearing rad nanno ooze.

Examples illustrating rules 1A and 1B and the resulting sediment names:

<u>% Clay</u>	<u>% Nannos</u>	=	
0- 5	75-95	=	Nanno ooze
10- 25	25-75	=	Clay-bearing nanno ooze
75- 95	5-25	=	Nanno-bearing clay
95-100	0- 5	=	Clay

II. Specific rules for calcareous and siliceous tests.

- A. Nannofossil is applied only to the calcareous tests of coccolithophorids, discoasters, etc.
- B. The term calcareous or siliceous, depending on skeletal composition, is applied where no attempt is made to distinguish fossils as to major subgroup. Thus, if no estimate is made, a mixture of radiolarians, diatoms, and silicoflagellates would be called a siliceous ooze. Where this distinction is made, the appropriate fossil name is used.
- C. Fossil tests are not qualified by a textural term unless very obviously redeposited.
- D. Abbreviations, as nanno for nannofossil, rad for radiolarian, etc. may be used in sediment name.

- E. The terms ooze, marl, and clay are used to designate >60%, 30 to 60%, and <30% carbonate, respectively (see Table 1).
- F. The term chalk is used to represent a compacted, semi-lithified ooze.
- G. Limestone is restricted to cemented calcareous rocks.

Table 1. Shales apply when sediment is fissile. Limestones must be cemented.

CaCO ₃ Content Percent	INDURATION		
	Unconsolidated	Semilithified	Lithified
0-30	clay	clay	claystone
30-60	marl	marl	marlstone
>60	ooze	chalk	limestone

III. Clastic sediments

- A. Clastic constituents, whether detrital, volcanic, biogenous or authigenic, are given a textural designation. When detrital¹ grains are the sole clastic constituents of a sediment, a simple textural term suffices for its name. The textural term can be preceded by a mineralogical term when this seems warranted. Such mineralogical terms are applied as per rules 1A and B.

B. Clastic volcanics

Redeposited pyroclastics also become a clastic component. They are recognized by the term volcanic and receive a

¹ Detrital = all clastic grains derived from the erosion of pre-existing rocks except for those of biogenous, authigenic, or volcanic origin.

textural term such as gravel, sand, silt, etc. It is particularly difficult at times to differentiate between volcanic sand (i.e., transported by tractive mechanisms) and crystal ash (i.e., direct outfall resulting from explosion of a volcano).

C. Clastic authigenic constituents

Where authigenic minerals are recognized as being a redeposited constituent, they are given a textural designation in addition to their mineral names.

IV. Volcanic and authigenic constituents

A. Volcanic constituents

Pyroclastics are given textural designations already established in the literature. Thus, volcanic breccia = >32 mm, volcanic lapilli = <32 mm to >4 mm, and volcanic ash = <4 mm. It is at times useful to further refine the textural designations by using such modifiers as coarse or fine. An ash wholly, or almost wholly, of glass shards is termed vitric ash.

B. Authigenic constituents

1. Authigenic minerals enter the sediment name in a fashion similar to that outlined under rules 1A and B. Normally, as with a fossil biocoenosis, the authigenic minerals are not given a textural designation and texture.
2. The terms ooze and chalk are applied to carbonate minerals of all types using the same rules that apply to biogenous constituents.

The lithologic symbols used in the core forms are shown in Figure 2.
Determinations of Shipboard Mineralogy and Lithology.

Smear Slides

Smear slides are the means of mineral identification on shipboard. Smear slide estimates of mineral abundances were based on the area of the smear slide covered by each component. Quantification was assigned using the following criteria:

Smear Slide descriptions

X-ray, grain size, and carbon-carbonate data

Many cores contain important minor lithologies as well as a basic lithology. The description of the basic lithology is so indicated in most cases; however, descriptive information for minor lithologies is included wherever possible.


A sample core form appears in Figure 3. This sample core form contains all legend and explanatory notes for understanding of the core forms.

Biostratigraphy

The biostratigraphy used for Leg 41 material follows the references in Table 2.

Table 2. Biostratigraphy

	Cenozoic	Mesozoic
Nannofossils	Martini, 1971	Thierstein, 1971
Foraminifera	Bolli, 1966 Banner and Blow, 1965	van Hinte, 1972
Radiolaria	Riedel and Sanfilippo (in press)	Riedel and Sanfilippo, 1974



V = Volcanic Ash
 F = Forams
 R = Rads
 D = Diatoms
 N = Nannos

Z = Zeolitic
 △ = Turbidite
 Fe = Ferruginous
 △ = Porcellanite
 ▲ = Chert
 A = Gypsum or Barite

Figure 2. Lithologic Symbols

Site	Hole	Core	Cored Interval:		LITHOLOGIC DESCRIPTION							
			AGE	NANNOFOSSIL ZONES FORAMINIFERA ZONES RADIODOLARIA ZONES	FOSSIL CHARACTER	FOSSIL ABUND.	PRES.	SECTION METERS	LITHOLOGY	DEFORIFICATION	LITHO SAMPLE	
				D = diatom, F = foraminifera, N = nannofossil, R = radiolaria, S = Silicoflagellate A = abundant, C = common, F = rare, T = trace, O = omitted G = good, M = moderate, P = poor				0				
								0.5				
								1.0				
								1				
								2				
								3				
								4				
								5				
								6				
		Core Catcher										
The lithology symbols are only to indicate the sediment types present and do not indicate relative amounts of each type.										Description of major and minor (if any) lithologies, color, deformation, and characteristics.		
SMEAR SLIDE - section, depth in section Abundances are qualitative follow the general scheme:										Description of major and minor (if any) lithologies, color, deformation, and characteristics.		
--- moderate disturbance, ~~~~severe disturbance, bx = drilling breccia, VOID = no material recovered Blank = minor to no disturbance										Description of major and minor (if any) lithologies, color, deformation, and characteristics.		
*Smear Slide depth in centimeters within the section C = chemistry, TS = thin section										Description of major and minor (if any) lithologies, color, deformation, and characteristics.		

Explanatory notes in Chapter 1

Figure 3. Sample Core Form and Legends

Drilling Deformation

Four degrees of drilling deformation were recognized and are noted by symbols on the sample core form (Figure 3). Slightly deformed cores exhibit a slight bending of bedding contacts; extreme bending defines moderate deformation. In highly deformed cores, injected bedding may be completely disrupted to produce a "drilling breccia." Watery intervals generally have lost any bedding characteristics originally available. Great care and considerable judgment must be used in determining whether sediment structural features are original or are artifacts introduced by the drilling and coring techniques.

Downhole Contamination

Downhole contamination is a serious problem. Hard objects (manganese nodules, chert, lithic fragments, and pebbles) are often washed or dragged downhole. They are commonly lodged in the top of cores or incorporated into the middle of cores at levels far below their proper stratigraphic position. Displaced manganese nodules can usually be detected. However, displaced chert, lithic fragments, and pebbles are more difficult to recognize.

1- 5% Rare (R)

5-25% Common (C)

25-75% Abundant (A)

>75% Dominant (D)

18

Specific mineral identification and quantification was attempted for sands, but for silts, and clays, only the textural categories were used.

Core Forms

The basic lithologic data are contained on core summary forms in both symbolic (Figure 2) and descriptive form. As far as possible the data are presented in the following order:

Sediment name

Color name and Munsell or GSA number

The reader is advised that colors recorded in core barrel summaries were determined during shipboard examination immediately after splitting the core sections. Experience with carbonate sediments has shown that many of the colors will fade or disappear with time after opening and storage. Colors particularly susceptible to rapid fading are purple, light and medium tints of blue, light bluish gray, dark greenish black, light tints of green, and pale tints of orange. These colors change to white or yellowish white or pale tan.

Composition

Structure(s)

REFERENCES

- Banner, F. T. and Blow, W. H., 1965. Progress in planktonic foraminiferal biostratigraphy of the Neogene: Nature, v. 208, no. 5016, p. 1164.

- Bolli, H., 1966. Zonation of Cretaceous to Pliocene marine sediments based on planktonic foraminifera: Bull. Assoc. Venezolana Geol. Mineral Petrol., v. 9, no. 1, p. 3.
- Krumbein, W. C. and Pettijohn, F. J., 1938. Manual of sedimentary petrography: New York, Appleton Century Co., Inc., p. 95-96.
- Martini, E., 1971. Standard Tertiary and Quaternary calcareous nannoplankton zonation: In Proc. 2nd Planktonic Conf., Roma, p. 739-785.
- Müller, G. and Gastner, M., 1971. The "karbonat-bombe", a simple device for the determination of the carbonate content in sediments, soils and other materials: N. Jb. Miner., H. 10, p. 466-469.
- Riedel, W. R. and Sanfilippo, A., 1974. Radiolaria from the Southern Indian Ocean, DSDP Leg 26: In Davis, T. A., Luyendyk, B. P., et al., Initial Reports of the Deep Sea Drilling Project, v. 26, Washington (U. S. Government Printing Office), p. 771-813.
- _____, in press. Stratigraphy and evolution of tropical Cenozoic Radiolaria: In Proc. 3rd Planktonic Conf. Kiel, Germany.
- Shepard, F. P., 1954. Nomenclature based on sand-silt-clay ratios: J. Sed. Petrol., v. 24, (3), p. 151-158.
- Thierstein, H. R., 1971. Tentative Lower Cretaceous calcareous nannoplankton zonation: Eclog. Geol. Helv., v. 64.
- Van Hinte, J. E., 1972. The Cretaceous time scale and planktonic foraminiferal zones: Koninkl. Nederl. Akad. vanwetenschappen, Series B, v. 75, no. 1, p. 8.
- Wentworth, C. K., 1922. A scale of grade and class terms of clastic sediments: J. Geol., v. 30, p. 377.

sample-distribution policy

Distribution of Deep Sea Drilling samples will be undertaken in order to (1) provide supplementary data for inclusion in the appropriate Initial Report to support *Glomar Challenger* scientists in achieving the scientific objectives of their particular cruise, and (2) provide individual investigators with material to conduct detailed studies beyond the scope of the Initial Reports.

The National Science Foundation has established a Sample Distribution Panel to advise on distribution of core material. This panel is chosen in accordance with usual Foundation practices, in a manner that will assure advice in the various disciplines leading to a complete and adequate study of the core and related materials. Funding for the proposed research is handled separately by the investigator, not through the Deep Sea Drilling Project.

Distribution of samples for contributions to Initial Reports

Any investigator who wishes to contribute a paper to a given volume of the Initial Reports may write to the Curator, Deep Sea Drilling Project, Scripps Institution of Oceanography, University of California at San Diego, La Jolla, 92037, requesting samples from a forthcoming cruise. The request should include the nature of the study, and type, size, number of samples, particular sampling techniques or equipment that might be required, and an estimate of the time required to complete the study. The requests will be reviewed by shipboard scientists, and, if they are deemed suitable and pertinent to the objectives of the leg, and shipboard workload permits, the requested samples will be taken during the cruise (provided, of course, material suitable to the investigation is obtained during the drilling). In the case of multiple requests to perform the same investigation, selection of investigator will be made by the shipboard scientific party.

Proposals should be of a scope appropriate to complete the sampling and study in time for publication in the Initial Reports. Studies deemed acceptable will be referred to the Curator who will, with the consent of the NSF Sample Distribution Panel, authorize distribution of the samples. The Sample Distribution Panel and the Deep Sea Drilling Project will strive to ensure a reasonable degree of continuity in the investigations among the various cruises, that the studies are pertinent to goals of the cruise, and that they are consistent with the publication policy for the Initial Reports. Subject to these same provisions, the shipboard scientific party may elect to have special studies of selected core samples of its recently completed cruise made by other investigators.

Investigations not completed in time for inclusion in the Initial Report may not be published in other journals until publication of the Initial Report for

which it was intended.

Distribution of samples for publication other than in Initial Reports

1. Researchers intending to request samples for studies beyond the scope of the Initial Reports should first obtain a sample request form from the Curator. Requests should specify the quantities and intervals of the core required, a statement of the proposed research, the possibility of returning residue to the Curator, the estimated time required to complete and publish the results, and the availability or need of funding and availability of equipment and space foreseen for the research.

In order to ensure that requests for highly desirable but limited samples can all be considered, approval of requests and distribution of samples will not be made prior to 12 months after date of completion of the cruise that collected the cores. Prior to publication of an Initial Report, requests for samples from a cruise can be based on the preliminary shipboard core logs. Copies of these logs will be kept on open file at Scripps and other designated institutions. The only exceptions will be for specific instances involving ephemeral properties.

Requests for samples from researchers in industrial laboratories will be handled in the same manner as those from academic organizations, and there will be the same obligation to publish results promptly. Requests from foreign scientists or organizations will also be considered. 2. The Curator has the responsibility for distributing samples, controlling quality of samples, and preserving core material. He also has the responsibility for maintaining a record of requests for samples that have been processed and filled indicating the investigator and subjects to be studied. This record will be available to investigators.

The distribution of samples will be made directly from the two repositories at Lamont-Doherty Geological Observatory and Scripps by the Curator or his designated representative.

3. (a) Samples up to 10 cc/m of core length can be automatically distributed by the Curator, Deep Sea Drilling Project or his authorized representative to any qualified investigator who requests them. The Curator will refrain from making automatic distribution of any parts of the cores which appear to be in particularly high demand, and any requests for these parts of the cores will be referred to the Sample Distribution Panel for review. Requests for samples from thin layers or important stratigraphic boundaries will generally require Panel review.

(b) All requests for samples in excess of 3(a) above will be referred to the Sample Distribution Panel.

(c) If, in the opinion of scientific investigators, certain properties they wish to study may deteriorate prior to the normal availability of the samples, such investigators may request that the normal waiting period not apply. All such requests

must be approved by the Sample Distribution Panel.

4. Samples will not be provided prior to assurance that funding for sample studies either exists or is not needed. However, neither formal approval of sample requests nor distribution of samples will be made until the appropriate time (Item 1). If a sample request is dependent, either wholly or in part, on proposed funding, the Curator will provide to the organization to whom the funding proposal has been submitted any information on the availability (or potential availability) of samples that it may request.

5. Investigators receiving samples are responsible for:

- i) promptly publishing significant results.
- ii) acknowledging, in publications, that samples were supplied through the assistance of the National Science Foundation.
- iii) submitting 4 copies of all reprints of published results to the Curator.
- iv) notifying the Curator of any work done on the samples that is additional to that stated in the original request for samples.

v) returning, in good condition, the remainders of samples after termination of research, if requested by the Curator.

6. Cores will be made available at repositories for investigators to examine and specify exact samples in such instances as this may be necessary for the scientific purposes of the sampling, subject to the limitations of 3 (a), (b), (c), and 5, above, and with the specific permission of the Curator or his delegate.

7. Cores of igneous and metamorphic rocks will also remain at the repositories where they will be available for observation and description and where selected samples may be taken for thin-section preparation and other work.

8. The Deep Sea Drilling Project routinely processes by computer most of the quantitative data presented in the Initial Reports. Space limits in the Initial Reports preclude detailed presentation of all such data. However, copies of the computer readout are available for those who wish the data for further analysis or as an aid in selecting samples.

Magnetics, seismic-reflection and bathymetric data collected under way by the *Glomar Challenger* will also be available for distribution 12 months after completion of the cruise.

Requests for these data may be made to the Coordinating Staff Geologist of the Deep Sea Drilling Project, at Scripps.

A charge will be made to recover the expenses of responding to individual requests. Estimated charges can be furnished before the request is processed, if required.

9. This policy has the approval of the National Science Foundation and is designed to help ensure that the greatest possible scientific benefit is gained from the materials obtained, and that samples will be made widely available to interested geologists.

(Slightly condensed from the official sample distribution policy of the Deep Sea Drilling Project.)

DEEP SEA DRILLING PROJECT

LEG 41 SITE 366 (HOLE 366A)

SITE SUMMARY SHEET

PRINCIPAL RESULTS:

Sierra Leone Rise

The sedimentary section on Sierra Leone Rise was continuously cored down to 850.5 meters and a complete Cenozoic record of pelagic sedimentation was recovered. The hole bottomed in upper Maestrichtian sediments and was abandoned due to plugging of the drill bit. The section consists of nanno oozes and marls grading downward to chalks and marls and then to limestones and marlstones. Chert and porcellanite were found in middle to lower Eocene. No hiatus sediments was found and all the biostratigraphic zones were sampled. The co-occurrence of different microfossil groups in most of the section makes it an ideal references section for Cenozoic zonation in tropical-subtropical latitudes. Since the latest Cretaceous Sierra Leone Rise was always in a deep-water pelagic environment, and well above the CCD. The rate of sedimentation appears rather constant with the lower values corresponding to times when hiatuses are observed in the basins and some other rises in the Atlantic. The highest rates of accumulation correspond to nonhiatus times. Dissolution and/or dilution cycles, present in most of the lower section up to the middle Miocene, are probably related to climatic variations. A major reflector observed at 0.5 seconds on the seismic profile corresponds to the youngest occurrence of middle Eocene porcellanites and chert. Another reflector, at about 0.9 seconds, but not corresponding to basement, was not reached so the age and nature of the oldest sediments remain unknown.

Date occupied	1140Z 22 February 1975
Date departed	1614Z 1 March 1975
Time on site	7 days, 4 hours, 34 minutes
Position: Latitude	05°40.7'N
Longitude	19°51.1'W
Water depth (sea level)	2853 corrected meters
Water depth (rig floor)	2863 corrected meters
Penetration	850.5 meters

LEG 41 SITE 366 (HOLE 366A)
SITE SUMMARY SHEET, con't.

Number of holes	2
Number of cores	55 at 366; 39 at 366A
Total length of cored section	850.5 meters
Total core recovered	582 meters
Percentage core recovery	68%
<u>Oldest Sediment Cored</u>	
Depth subbottom	950.5 meters
Nature	Limestone
Age	Maestrichtian

Site	366	Core Interval	119.0-128.5 m	LITHOLOGIC DESCRIPTION		LITHO. SAMPLE
				LITHOLOGY	DEFORMATION	
AGE						
ZONES						
CHARACTER						
FOSILS						
NANNO						
RAMS						
FORAMS						
MAMS						
ABUND.						
PRE.S.						
SECTION	0					
METERS	0.5					
1	1.0					
2	80					
3						
4						
5						
6						
Core						
Catcher						
CC						
F						
C						
P						
N						
R						
Globorotalia acostaeensis-Globigerina neopenthes Zones NN9-N11						
Discocaster hamatus-Discocaster quadrangularis NNG-N11						
UPPER MIOCENE-UPPER MIDDLE MIOCENE						
10R 7/4						
10R 8/2						
10R 8/2						
10R 8/2						
10R 8/2						
10R 8/2						
135						



Site 366		Core 8		Cover Interval: 394.5-404.0 m	
AGE	ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	LITHO. DESCRIPTION
			0		
			0.5	VOID	
			1	1.0	
			2		
			3		
			4		
			65		
			72		
			103		
			107		
			111		
			115		
			119		
			123		
			127		
			131		
			135		
			139		
			143		
			147		
			151		
			155		
			159		
			163		
			167		
			171		
			175		
			179		
			183		
			187		
			191		
			195		
			199		
			203		
			207		
			211		
			215		
			219		
			223		
			227		
			231		
			235		
			239		
			243		
			247		
			251		
			255		
			259		
			263		
			267		
			271		
			275		
			279		
			283		
			287		
			291		
			295		
			299		
			303		
			307		
			311		
			315		
			319		
			323		
			327		
			331		
			335		
			339		
			343		
			347		
			351		
			355		
			359		
			363		
			367		
			371		
			375		
			379		
			383		
			387		
			391		
			395		
			399		
			403		
			407		
			411		
			415		
			419		
			423		
			427		
			431		
			435		
			439		
			443		
			447		
			451		
			455		
			459		
			463		
			467		
			471		
			475		
			479		
			483		
			487		
			491		
			495		
			499		
			503		
			507		
			511		
			515		
			519		
			523		
			527		
			531		
			535		
			539		
			543		
			547		
			551		
			555		
			559		
			563		
			567		
			571		
			575		
			579		
			583		
			587		
			591		
			595		
			599		
			603		
			607		
			611		
			615		
			619		
			623		
			627		
			631		
			635		
			639		
			643		
			647		
			651		
			655		
			659		
			663		
			667		
			671		
			675		
			679		
			683		
			687		
			691		
			695		
			699		
			703		
			707		
			711		
			715		
			719		
			723		
			727		
			731		
			735		
			739		
			743		
			747		
			751		
			755		
			759		
			763		
			767		
			771		
			775		
			779		
			783		
			787		
			791		
			795		
			799		
			803		
			807		
			811		
			815		
			819		
			823		
			827		
			831		
			835		
			839		
			843		
			847		
			851		
			855		
			859		
			863		
			867		
			871		
			875		
			879		
			883		
			887		
			891		
			895		
			899		
			903		
			907		
			911		
			915		
			919		
			923		
			927		
			931		
			935		
			939		
			943		
			947		
			951		
			955		
			959		
			963		
			967		
			971		
			975		
			979		
			983		
			987		
			991		
			995		
			999		
			1003		
			1007		
			1011		
			1015		
			1019		
			1023		
			1027		
			1031		
			1035		
			1039		
			1043		
			1047		
			1051		
			1055		
			1059		
			1063		
			1067		
			1071		
			1075		
			1079		
			1083		
			1087		
			1091		
			1095		
			1099		
			1103		
			1107		
			1111		
			1115		
			1119		
			1123		
			1127		
			1131		
			1135		
			1139		
			1143		
			1147		
			1151		
			1155		
			1159		
			1163		
			1167		
			1171		
			1175		
			1179		
			1183		
			1187		
			1191		
			1195		
			1199		
			1203		
			1207		
			1211		
			1215		
			1219		
			1223		
			1227		
			1231		
			1235		
			1239		
			1243		
			1247		
			1251		
			1255		
			1259		
			1263		
			1267		
			1271		
			1275		
			1279		
			1283		
			1287		
			1291		
			1295		
			1299		
			1303		
			1307		
			1311		
			1315		
			1319		
			1323		
			1327		
			1331		
			1335		
			1339		
			1343		
			1347		
			1351		
			1355		
			1359		
			1363		
			1367		
			1371		
			1375		
			1379		
			1383		
			1387		
			1391		
			1395		
			1399		
			1403		
			1407		
			1411		
			1415		
			1419		
			1423		
			1427		
			1431		
			1435		
			1439		
			1443		
			1447		
			1451		
			1455		
			1459		
			1463		
			1467		
			1471		
			1475		
			1479		
			1483		
			1487		
			1491		
			1495		
			1499		
			1503		
			1507		
			1511		
			1515		
			1519		
			1523		
			1527		
			1531		
			1535		
			1539		

Site	Core	Interval	Lithological Description																						
			Fossil Character		AEC		Metres		Section		Lithology		Litho. Sample												
366	7	385.5-394.5 m	ZONES	Fossils	Rods	Rods	Prds.	Absnd.	0	1	1.0	1.0	100	SS at 1-100	Forams	Nannos	Rads	Sponge spicules	R	R-C	R	Core Catcher Chalk			
			FORAMS											SS at 3-22	(minor lithology)										
			MICRANOMS											PELAGIC CLAY	R-C	Diatoms									
			ZOOPLITES											SS at 4-120	(dominant lithology)										
			FORAMS											FORAMS	R	Diatoms	R								
			NRAMS											NANOMS	D	Rads	R								
			NRAMOS											NRAMOS	R										
			THELOCYRTIDS											THELOCYRTIDS	?										
			GLIOBLIGERITA											GLIOBLIGERITA	?										
			LEPTURINAE											LEPTURINAE	?										
			LOWER OLIGOCENE											Tuberosa		Nanoplankton		Diatoms		Diatoms		CC			
			5		120		103		22		100		SS at 5-90		(dominant lithology)		Diatoms		Diatoms		C				
			4		120		103		22		100		SS at CC		(?)		Diatoms		Diatoms		C				
			3		120		103		22		100		ZOOPLITES (?)		?		Diatoms		Diatoms		C				
			2		120		103		22		100		NANOMS		A		Sponge spicules		Sponge spicules		R				
			1		120		103		22		100		NANOMS		A		Rads		Rads		R				
			0		120		103		22		100		NANOMS		D		Rads		Rads		R				

MIDDLE EOCENE		LITHOLOGY		DEFORMATION		LITHO. SAMPLE		LITHOLOGIC DESCRIPTION	
AGE	ZONES	FOSIL	CHARACTER	METERS	SECTION	METERS	SECTION		
FORAMS	TRUNCOCYLINDRIDIENS ROTH ET PIA	N	R	0					INTERBEDDED PORCELLANITE AND NANO CHALK IN CYCLES (>25 cm)
FORAMS	DISCOASPIRIDES TANZ MOLLET NP15-NP16	F	C	0.5					NANO CHALK, light greenish gray (5GY 8/1), well-indurated but not cemented, severe drilling disturbance, faintish and burrowed.
NANNO	DISCOASPIRIDES TANZ MOLLET NP15-NP16	P	P	1					PORCELLANITE, light gray (N7) becoming cherry gray (5Y 6/1).
				1.0					SS at 1-40 (dominant lithology)
									Recrystallized calcite A Namnos A
									SS at 1-80 (dominant lithology)
									Recrystallized calcite C Forams R
									Carbonate Bomb: 1-103 to 104 cm = 83%

31

Site 366 Hole		Core 22 Cored Interval : 527.5-537.0 m			
ZONES	FOSSTIL CHARACTER	AGE	FOAMS	MATERIALS	
SECTIION	LITHOLOGY	METERS	LITOTHO. SAMPLE	DEFORMATION	LITHOLOGIC DESCRIPTION
0	PRES.	0			INTERBEDDED NANNO CHALK, SILICIFIED LIMESTONE, AND CHERI IN CYCLES (~25 cm) NANNO CHALK, light greenish gray (SGY 8/1), well-indurated, laminated. Grades into SILICIFIED LIMESTONE.
0.5	AIRBUD.				CHERI, light gray (N7), shows burrows, laminated.
1	FOSSTIL	1.0			SS at CC Recrystallized calcite
	PRES.				Forams Nannos
	AIRBUD.				C
	FOSSTIL				C
	PRES.				R
2					Carbonate Bomb: 1-49 to 50 cm = 83%
					C
					CC
					Core Catcher





37

Site 366 Hole		Core 39	Cored Interval: 689.0-698.5 m	LITHOLOGIC DESCRIPTION						
AGE	ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY		DEFORMATION		LITHO. SAMPLE		LOWER EOCENE
				SECTIION	PRES.	ABUND.	FOSIL	NANNO	RAMS	
			0	VOID						
			1	0.5						
			1	1.0						
			2							
			3							
			4							
			5							
			6							
										Core Catcher


Site 366 Hole		Core 40	Cored Interval: 698.5-708.0 m	LITHOLOGIC DESCRIPTION						
AGE	ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY		DEFORMATION		LITHO. SAMPLE		LOWER EOCENE
				SECTIION	PRES.	ABUND.	FOSIL	NANNO	RAMS	
			0	VOID						
			1	0.5						
			1	1.0						
			2							
			3							
			4							
			5							
			6							
										Core Catcher

Site 366 Hole		Core 42	Core interval: 717.5-727.0 m
ZONES	CHARACER	FUSSIL	PRBS.
AG6			
FORMAMS	RADS		
NANNOS	NANOS		
Forams	Rods		
ABUND.			
FUSSIL			
SECTIION	METERS	LITHOLOGY	DEFORMATION
0	0.5-	ANGILLACEOUS LIMESTONE, greenish gray (N7) to light greenish gray (56y 8/1), finely laminated, burrowed. Wimmed layers of CALCARENITE, occur at 3-100, 3-27, 4-18, 4-145, 5-60, 5-105, 5-130, 6-45, 6-90, 6-103, 6-110, 6-20, 6-140.	LITHO. SAMPLE
1	1.0-	SS at 3-90 (dominant lithology) Clay Forams Nanios SS at 3-127 (minor lithology) Recrystallized calcite A Nanios SS at CC Recrystallized calcite A	SS at 3-90 (dominant lithology) Clay Forams Nanios SS at 3-127 (minor lithology) Recrystallized calcite A Nanios SS at CC Recrystallized calcite A
2			Carbonate Bomb: 3-123 to 124 cm = 71%
3			90
4			127
5			
6			
			Core Catcher



Site 366 Hole		Core 47	Cored Interval: 765.0-774.5 m
		LITHO. SAMPLE	DEFORMATION
		LITHOLOGIC DESCRIPTION	
DEFORMATION	LITHO. SAMPLE		
LITHOLOGY	LITHOLOGIC DESCRIPTION		
METERS			
SECTION			
0	0		
0.5	SLILOCIOUS LIMESTONE, light gray (N7), finely laminated, burrowed.		
1	SS at 3-60 (dominant lithology) Clay C Recrystallized Calcite crystals R calcite A Forams R Namus A Carbonate Bomb: 3-43 to 44 cm = 65%		
1.0			
2			
3			
4			
5			
6			
6.0			
AGE	FOSIL CHARACTER	PRBS.	
ZONES	FORAMS	ABUND.	
	NANNO		
	RADS		
	FOSIL		
	CHARACT.		
	PRBS.		
	SECTION		
	METERS		
	LITHOLOGY		
	LITHO. SAMPLE		
	DEFORMATION		

Site 366 Hole		Core 48 Cored Interval: 774.5-784.0 m			
ZONES	AGE	FORMS	RADS	FOSIL	ABUND.
CHARACTER	FOSSIL	PRBS.	PRES.	SECTION	METERS
Heijolitithus pseudomenardi P4					0
Globigerinata pseudomenardi NP8					0.5
NANNO					1
RAD5					1.0
FORMS					2
MANNS					3
MANNS					4
					5
					6
					7
					8
					9
					10
					11
					12
					13
					14
					15
					16
					17
					18
					19
					20
					21
					22
					23
					24
					25
					26
					27
					28
					29
					30
					31
					32
					33
					34
					35
					36
					37
					38
					39
					40
					41
					42
					43
					44
					45
					46
					47
					48
					49
					50
					51
					52
					53
					54
					55
					56
					57
					58
					59
					60
					61
					62
					63
					64
					65
					66
					67
					68
					69
					70
					71
					72
					73
					74
					75
					76
					77
					78
					79
					80
					81
					82
					83
					84
					85
					86
					87
					88
					89
					90
					91
					92
					93
					94
					95
					96
					97
					98
					99
					100
					101
					102
					103
					104
					105
					106
					107
					108
					109
					110
					111
					112
					113
					114
					115
					116
					117
					118
					119
					120
					121
					122
					123
					124
					125
					126
					127
					128
					129
					130
					131
					132
					133
					134
					135
					136
					137
					138
					139
					140
					141
					142
					143
					144
					145
					146
					147
					148
					149
					150
					151
					152
					153
					154
					155
					156
					157
					158
					159
					160
					161
					162
					163
					164
					165
					166
					167
					168
					169
					170
					171
					172
					173
					174
					175
					176
					177
					178
					179
					180
					181
					182
					183
					184
					185
					186
					187
					188
					189
					190
					191
					192
					193
					194
					195
					196
					197
					198
					199
					200
					201
					202
					203
					204
					205
					206
					207
					208
					209
					210
					211
					212
					213
					214
					215
					216
					217
					218
					219
					220
					221
					222
					223
					224
					225
					226
					227
					228
					229
					230
					231
					232
					233
					234
					235
					236
					237
					238
					239
					240
					241
					242
					243
					244
					245
					246
					247
					248
					249
					250
					251
					252
					253
					254
					255
					256
					257
					258
					259
					260
					261
					262
					263
					264
					265
					266
					267
					268
					269
					270
					271
					272
					273
					274
					275
					276
					277
					278
					279
					280
					281
					282
					283
					284
					285
					286
					287
					288
					289
					290
					291
					292
					293
					294
					295
					296
					297
					298
					299
					300
					301
					302
					303
					304
					305
					306
					307
					308
					309
					310
					311
					312
					313
					314
					315
					316
					317
					318
					319
					320
					321
					322
					323
					324
					325
					326
					327
					328
					329
					330
					331
					332
					333
					334
					335
					336
					337
					338
					339
					340
					341
					342
					343
					344
					345
					346
					347
					348
					349
					350
					351
					352
					353
					354
					355
					356
					357
					358
					359
					360
					361
					362
					363
					364
					365
					366
					367
					368
					369
</					



Site 366		Core 52		Cored Interval: 812.5-822.0 m			
AGE	ZONES	FOSSIL CHARACTER	FOSSIL	SECTION METERS	LITHOLOGY	DEFORMATION	LITHO. SAMPLE
PALEOCENE, DANIAN S-S.	Globorotalia trituberculata tenus NP2	MAMS	F A M R N R P	0	PRBS. AGUND.		
							MARLSTONE, light olive gray (SY 6/1).

Site 366		Hole	Core 55	Cored Interval: 841.0-850.5 m									
AGES	ZONES	FOSIL CHARACTER	SECTIION METERS	LITHOLOGY	LITHOLOGIC DESCRIPTION								
					F	A	M	N	p-N	CORE CATCHER	DEFORMATION	LITHO. SAMPLE	DEFNATION
GRETACEOUS, MESTRICHITIAN	Globorotalites quadratus	ABUND.	0	SS at CC							ONLY SCRAPINGS OFF OF CORE CATCHER MARLSTONE, light olive gray (5Y 6/1).		
	FORMAMS	F		Clay							Fe/Mn		
	RADS	N		Recrystallized							micronodules	R	
	MANNOS	F		calcite							Calcite crystals	R	
		A		Nannos							Fish debris	R	

Site	366	Hole A	Core 1		Cored Interval: 0.0-6.0 m
			AGE	LITHOLOGY	
ZONES		FOSIL CHARACTER		METRES	LITHO. & SAMPLE
FORAMS		ABUND.		0	DEFORMATION
NANNO		PRDS.		0.5	
				1	
				1.0	
				1.5	
				2	
				3	
				4	
				5	
				6	
				7	
				8	
				9	
				10	
				11	
				12	
				13	
				14	
				15	
				16	
				17	
				18	
				19	
				20	
				21	
				22	
				23	
				24	
				25	
				26	
				27	
				28	
				29	
				30	
				31	
				32	
				33	
				34	
				35	
				36	
				37	
				38	
				39	
				40	
				41	
				42	
				43	
				44	
				45	
				46	
				47	
				48	
				49	
				50	
				51	
				52	
				53	
				54	
				55	
				56	
				57	
				58	
				59	
				60	
				61	
				62	
				63	
				64	
				65	
				66	
				67	
				68	
				69	
				70	
				71	
				72	
				73	
				74	
				75	
				76	
				77	
				78	
				79	
				80	
				81	
				82	
				83	
				84	
				85	
				86	
				87	
				88	
				89	
				90	
				91	
				92	
				93	
				94	
				95	
				96	
				97	
				98	
				99	
				100	
				101	
				102	
				103	
				104	
				105	
				106	
				107	
				108	
				109	
				110	
				111	
				112	
				113	
				114	
				115	
				116	
				117	
				118	
				119	
				120	
				121	
				122	
				123	
				124	
				125	
				126	
				127	
				128	
				129	
				130	
				131	
				132	
				133	
				134	
				135	
				136	
				137	
				138	
				139	
				140	
				141	
				142	
				143	
				144	
				145	
				146	
				147	
				148	
				149	
				150	
				151	
				152	
				153	
				154	
				155	
				156	
				157	
				158	
				159	
				160	
				161	
				162	
				163	
				164	
				165	
				166	
				167	
				168	
				169	
				170	
				171	
				172	
				173	
				174	
				175	
				176	
				177	
				178	
				179	
				180	
				181	
				182	
				183	
				184	
				185	
				186	
				187	
				188	
				189	
				190	
				191	
				192	
				193	
				194	
				195	
				196	
				197	
				198	
				199	
				200	
				201	
				202	
				203	
				204	
				205	
				206	
				207	
				208	
				209	
				210	
				211	
				212	
				213	
				214	
				215	
				216	
				217	
				218	
				219	
				220	
				221	
				222	
				223	
				224	
				225	
				226	
				227	
				228	
				229	
				230	
				231	
				232	
				233	
				234	
				235	
				236	
				237	
				238	
				239	
				240	
				241	
				242	
				243	
				244	
				245	
				246	
				247	
				248	
				249	
				250	
				251	
				252	
				253	
				254	
				255	
				256	
				257	
				258	
				259	
				260	
				261	
				262	
				263	
				264	
				265	
				266	
				267	
				268	
				269	
				270	
				271	
				272	
				273	
				274	
				275	
				276	
				277	
				278	
				279	
				280	
				281	
				282	
				283	
				284	
				285	
				286	
				287	
				288	
				289	
				290	
				291	
				292	
				293	
				294	
				295	
				296	
				297	
				298	
				299	
				300	
				301	
				302	
				303	
				304	
				305	
				306	
				307	
				308	
				309	
				310	
				311	
				312	
				313	
				314	
				315	
				316	
				317	
				318	
				319	
				320	
				321	
				322	
				323	
				324	
				325	
				326	
				327	
				328	
				329	
				330	
				331	
				332	
				333	
				334	
				335	
				336	
				337	
				338	
				339	
				340	
				341	
				342	
				343	
				344	
				345	
				346	
				347	
				348	
				349	
				350	
				351	
				352	
				353	
				354	
				355	
				356	
				357	
				358	
				359	
				360	
				361	
				362	
				363	
				364	
				365	
				366	
				367	
				368	
				369	
				370	

46

Site 366		Core A	Core 3	Cored Interval: 15.5-25.0 m
AGES	ZONES	FUSSIL	LITHOLOGY	LITHO. SAMPLE
FORAMS	FORAMS	FOSSIL CHARACTER	DEFORMATION	LITHO. SAMPLE
NANNO	NANNO	ABUND.		
FASS	FASS	RELS.		
SECTIION	METERS			
	0		VOID	
	1			
	1.0			
	0.5			
	1			
	2			
	3			
	4			
	5			
	6			
	7			
	8			
	9			
	10			
	11			
	12			
	13			
	14			
	15			
	16			
	17			
	18			
	19			
	20			
	21			
	22			
	23			
	24			
	25			
	26			
	27			
	28			
	29			
	30			
	31			
	32			
	33			
	34			
	35			
	36			
	37			
	38			
	39			
	40			
	41			
	42			
	43			
	44			
	45			
	46			
	47			
	48			
	49			
	50			
	51			
	52			
	53			
	54			
	55			
	56			
	57			
	58			
	59			
	60			
	61			
	62			
	63			
	64			
	65			
	66			
	67			
	68			
	69			
	70			
	71			
	72			
	73			
	74			
	75			
	76			
	77			
	78			
	79			
	80			
	81			
	82			
	83			
	84			
	85			
	86			
	87			
	88			
	89			
	90			
	91			
	92			
	93			
	94			
	95			
	96			
	97			
	98			
	99			
	100			
	101			
	102			
	103			
	104			
	105			
	106			
	107			
	108			
	109			
	110			
	111			
	112			
	113			
	114			
	115			
	116			
	117			
	118			
	119			
	120			
	121			
	122			
	123			
	124			
	125			
	126			
	127			
	128			
	129			
	130			
	131			
	132			
	133			
	134			
	135			
	136			
	137			
	138			
	139			
	140			
	141			
	142			
	143			
	144			
	145			
	146			
	147			
	148			
	149			
	150			
	151			
	152			
	153			
	154			
	155			
	156			
	157			
	158			
	159			
	160			
	161			
	162			
	163			
	164			
	165			
	166			
	167			
	168			
	169			
	170			
	171			
	172			
	173			
	174			
	175			
	176			
	177			
	178			
	179			
	180			
	181			
	182			
	183			
	184			
	185			
	186			
	187			
	188			
	189			
	190			
	191			
	192			
	193			
	194			
	195			
	196			
	197			
	198			
	199			
	200			
	201			
	202			
	203			
	204			
	205			
	206			
	207			
	208			
	209			
	210			
	211			
	212			
	213			
	214			
	215			
	216			
	217			
	218			
	219			
	220			
	221			
	222			
	223			
	224			
	225			
	226			
	227			
	228			
	229			
	230			
	231			
	232			
	233			
	234			
	235			
	236			
	237			
	238			
	239			
	240			
	241			
	242			
	243			
	244			
	245			
	246			
	247			
	248			
	249			
	250			
	251			
	252			
	253			
	254			
	255			
	256			
	257			
	258			
	259			
	260			
	261			
	262			
	263			
	264			
	265			
	266			
	267			
	268			
	269			
	270			
	271			
	272			
	273			
	274			
	275			
	276			
	277			
	278			
	279			
	280			
	281			
	282			
	283			
	284			
	285			
	286			
	287			
	288			
	289			
	290			
	291			
	292			
	293			
	294			
	295			
	296			
	297			
	298			
	299			
	300			
	301			
	302			
	303			
	304			
	305			
	306			
	307			
	308			
	309			
	310			
	311			
	312			
	313			
	314			
	315			
	316			
	317			
	318			
	319			
	320			
	321			
	322			
	323			
	324			
	325			
	326			
	327			
	328			
	329			
	330			
	331			
	332			
	333			
	334			
	335			
	336			
	337			
	338			
	339			
	340			
	341			
	342			
	343			
	344			
	345			
	346			
	347			
	348			
	349			
	350			
	351			
	352			
	353			
	354			
	355			
	356			
	357			
	358			
	359			
	360			
	361			
	362			
	363			
	364			
	365			
	366			
	367			
	368			
	369			
	370			
	371			
	372			
	373			
	374			
	375			
	376			
	377			
	378			
	379			
	380			
	381			
	382			
	383			
	384			
	385			
	386			
	387			
	388			
	389			
	390			
	391			
	392			
	393			
	394			
	395			
	396			
	397			
	398			
	399			
	400			
	401			
	402			
	403			
	404			
	405			
	406			
	407			
	408			
	409			
	410			
	411			
	412			
	413			
	414			
	415			
	416			
	417			
	418			
	419			
	420			
	421			

Site 366 Hole A		Core 5		Corred Interval : 34.5-44.0 m	
ZONES	CHARACTER	FOSIL	CHARACTER	FOSIL	DEFORMATION
AGE	RAMS	NANNO	RAMS	NANNO	LITHO. SAMPLE
0	PRES.	0	0	N	LITHOLOGY
0.5	ABUND.	0.5	1	N	MARL, varicolored because of drilling disturbance, from white (W) to greenish gray (SG 6/1) to dark yellowish green (SGY 6/1), soft, intense drilling disturbance, black mottling occurs at 3.0 to 7.0 cm; 12-20 to 30 cm; 5.0 to 35 cm; 5-40 to 90 cm.
1	Fossils	1	1.0	N	SS at CC (dominant lithology)
2	Nannons	2	2	N	Clay
3	Forams	3	3	N	Forams
4	Nannons	4	4	N	Nannons
5	Forams	5	5	N	Fish debris
6	Nannons	6	6	N	Carbonate Bomb: 3.76 to 77 cm = 66*
					Core Catcher

Site 366 Hole A		Core 11 Cored Interval: 91.5-101.0 m		LITHOLOGIC DESCRIPTION	
ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY	DEFORMATION	LITHO. SAMPLE
FORAMS	ABUND.	0	VOID		
NANNO	FOSSIL	0.5			
RADS	PRBS.	1			
NANNO	ABUND.	1.0			
RADS	PRBS.	2			
NANNO	ABUND.	3			
RADS	PRBS.	4			
NANNO	ABUND.	5			
RADS	PRBS.	6			
NANNO	ABUND.	100			
RADS	PRBS.	122			
NANNO	ABUND.	CC			

Globorotalia marginatae N19 base-N18 Ceratotilus tricarinatus-Reticulofenestrina pseudolumifera NNT2-NNT5 Globigerina plesiotumida N17 Discaster quinqueramus N111

UPPER Miocene

Site 366 Hole A		Core 12 Cored Interval: 101.0-110.5 m		LITHOLOGIC DESCRIPTION	
ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY	DEFORMATION	LITHO. SAMPLE
FORAMS	ABUND.	0	VOID		
NANNO	FOSSIL	0.5			
RADS	PRBS.	1			
NANNO	ABUND.	1.0			
RADS	PRBS.	2			
NANNO	ABUND.	3			
RADS	PRBS.	4			
NANNO	ABUND.	5			
RADS	PRBS.	6			
NANNO	ABUND.	100			
RADS	PRBS.	122			
NANNO	ABUND.	CC			

NANNO Ooze: yellowish gray (5Y 8/1) with white (N9) mottles. Soupy, intense drilling disturbance. Section 4 has black streaks throughout. Color gradually changes to very pale brown (10R 7/3) with white (N9) mottles by Section 4.

SS at CC Clay Forams Fe/Mn Rads Fish debris

SS at CC Clay Forams Nannos Rads Fish debris

Carbonate Bomb: 3-54 to 55 cm = 83%

NANNO Ooze: white (N9) with mottles of yellowish gray (5Y 8/1). Soupy in upper section, soft in lower 4 sections, intense drilling disturbance. Black streaks throughout Sections 2, 3 and 6.

SS at 6-100 (dominant lithology) Clay Forams Fe/Mn Rads Fish debris

SS at CC Clay Forams Nannos Rads Fish debris

Carbonate Bomb: 3-110 to 111 cm = 72%

Site 366 Hole A Core 13 Cored Interval: 110.5-120.0 m									
Globorotalia plesiostomula NT7 Discocaster quinqueramus NT1									
UPPER Miocene									
ZONES	FOSIL CHARACTER	LITHOLOGY	METERS	SECTION	PRBS.	FOSILL	RADS.	NANNO.	AGE
FORAMS	NANNO	SS at CC Clay Forams	0	0.5	A	Nannos C	A	R	
			1	1.0					
			2	10.0					
			3	10.0					
			4	10.0					
			5	10.0					
			6	10.0					
									Core Catcher
									CC

Site 366 Hole A Core 14 Cored Interval: 120.0-129.5 m									
Globorotalia acostaeensis NT6 Discocaster quinqueramus NT1									
UPPER Miocene									
ZONES	FOSIL CHARACTER	LITHOLOGY	METERS	SECTION	PRBS.	FOSILL	RADS.	NANNO.	AGE
FORAMS	NANNO	SS at CC Clay Forams	0	0.5	A	Nannos R	A	R	
			1	1.0					
			2	10.0					
			3	10.0					
			4	10.0					
			5	10.0					
			6	10.0					
									Core Catcher
									CC

Site 366		Core 16 Hole A		Core Interval: 139.0-148.5 m	
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	MATERIAL	LITHO. SAMPLE
					LITHOLOGIC DESCRIPTION
					DRILLING BRECCIA OF PELAGIC CLAY/MARL AND NANNO Ooze IN SECTION 1 AND 2. Pale grayish orange (DOR 8/4) to yellowish brown (DOR 5/3). Soft to stiff, drilling breccia.
					At 3-102 cm: NANNO MARL, greenish gray (SGY 6/1) to light greenish gray (SGY 8/1) w/ mottles of light brown (DOR 7/4), stiff, intense drilling disturbance.
					Section 5: Cyclic alternations of the above NANNO MARL and a NANNO Ooze, white (DOR 8/1), stiff, slight drilling disturbance. The nanno marls are generally grayish orange (DOR 7/4).
					Each cycle is about 20 cm thick but they gradually shorten to about 10 cm thick the last third of Section 5.
					SS at 3-140 (minor lithology) Clay A Forams Nanom R Radls Fish debris R
					SS at 5-122 (minor lithology) Clay D Fe/Mn Forams R Nanos Radls R Fish
					SS at CC (dominant lithology) Clay A Fe/Mn Amorph SiO ₂ R Forams Nanom A Radls Radls R Fish
					Carbonate Bomb: 3-140 to 141 cm = 60%
0	0	VOID	bx		
1	1	0.5			
2	2	1.0			
3	3	1			
4	4				
5	5				
					F-C P A G F N R N Core Catcher

52

Site 366 Hole A		Core 20 Cored Interval : 177.0-186.5 m			
ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY	LITHOLOGIC DESCRIPTION	
				LITHO. & SAMPLE	DEFORMATION
AE6		0	VOID	FORAM NANNO CHALK, light greenish gray (SY 8/1), stiff, slight drilling disturbance. Mottled with yellowish gray (SY 8/1).	R
FORAMS	NANNO.	0.5		SS at CC	A
RADLS.	ABUND.	1.0		Clay	Fe/Mn
				Forams	Nanno
				Rads	Sponge spicules
					R
				Carbonate Bomb.	3-104-105 cm = 77*
Globigerinata Statiferi-Globigerinata dissimilis N6-N5 Sericostasea trilobata-Serpenotithus belemnos NN2-NN3					
Stichoceras woffitti					
AEG					
LOWER MIOCENE					
F N F P-A Core					
F F A G Catcher					
R F E					
CC					

Site 366 Hole A		Core 24		Cored Interval: 215.0-224.5 m		Core 26		Cored Interval: 234.0-243.5 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									
ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER
FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.
RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	FOSIL	DEFORMATTION	RDPS	RADPS	FOSIL	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
LITHOLOGIC DESCRIPTION									
LITHOLOGIC DESCRIPTION									
CLAYEY NANNO CHALK, light greenish gray (SG 8/1), stiff, slight to severe drilling disturbance, burrowed. Thin (10 cm) interbeds of slightly darker green occur at 2-70 to 80 cm; 3-67 to 54 cm; 3-85 to 100 cm; scattered throughout Section 4 and at 5-20 to 30 cm.									
These slightly darker green layers appear to have less carbonate and may be part of the previous dissolution cycle sequence.									
SS at 4-65 (dominant lithology)									
Clay Forams Rads									
SS at CC									
Clay Forams Rads									
Carbonate Bomb: 3-113 to 114 cm = 80%									
Carbonate Bomb: 3-113 to 114 cm = 80%									
LOWER MIocene									
GLOBALIGERINATA DISSIMILITIS N5									
CYRTOCARPAS TESTICRIPPA									
DISCOASTER DRUGGET N2									
LYCHMECHANOMA ELONGATA									
Site 366 Hole A		Core 25		Cored Interval: 224.5-234.0 m		Core 26		Cored Interval: 234.0-243.5 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									
ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER
FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.
RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	FOSIL	DEFORMATTION	RDPS	RADPS	FOSIL	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
LITHOLOGIC DESCRIPTION									
LITHOLOGIC DESCRIPTION									
FORAN-DIATOR-BEARING CLAYEY NANNO CHALK, light greenish gray (SG 8/1), stiff, severe drilling disturbance, some darker green burrowed segments. <i>Ethmiodiscus rex</i> common.									
SS at CC									
Clay Forams Rads									
CC									
LOWER MIocene									
GLOBALIGERINATA DISSIMILITIS N5									
CYRTOCARPAS TESTICRIPPA									
DISCOASTER DRUGGET N2									
LYCHMECHANOMA ELONGATA									
Site 366 Hole A		Core 24		Cored Interval: 215.0-224.5 m		Core 25		Cored Interval: 224.5-234.0 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									
ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER
FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.
RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	FOSIL	DEFORMATTION	RDPS	RADPS	FOSIL	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
LITHOLOGIC DESCRIPTION									
LITHOLOGIC DESCRIPTION									
FORAN-DIATOR-BEARING CLAYEY NANNO CHALK, light greenish gray (SG 8/1), stiff, severe drilling disturbance, some darker green burrowed segments. <i>Ethmiodiscus rex</i> common.									
SS at CC									
Clay Forams Rads									
CC									
LOWER MIocene									
GLOBALIGERINATA DISSIMILITIS N5									
CYRTOCARPAS TESTICRIPPA									
DISCOASTER DRUGGET N2									
LYCHMECHANOMA ELONGATA									
Site 366 Hole A		Core 24		Cored Interval: 215.0-224.5 m		Core 25		Cored Interval: 224.5-234.0 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									
ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER
FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.
RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	FOSIL	DEFORMATTION	RDPS	RADPS	FOSIL	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
LITHOLOGIC DESCRIPTION									
LITHOLOGIC DESCRIPTION									
FORAN-DIATOR-BEARING CLAYEY NANNO CHALK, light greenish gray (SG 8/1), stiff, severe drilling disturbance, some darker green burrowed segments. <i>Ethmiodiscus rex</i> common.									
SS at CC									
Clay Forams Rads									
CC									
LOWER MIocene									
GLOBALIGERINATA DISSIMILITIS N5									
CYRTOCARPAS TESTICRIPPA									
DISCOASTER DRUGGET N2									
LYCHMECHANOMA ELONGATA									
Site 366 Hole A		Core 24		Cored Interval: 215.0-224.5 m		Core 25		Cored Interval: 224.5-234.0 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									
ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER	METERS	LITHOLOGY	ZONES	CHARACTER
FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.	PRES.	DEFORMATION	FORAMS	NANNO.
RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS	ABUND.	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	FOSIL	DEFORMATTION	RDPS	RADPS	FOSIL	LITHO-SAMPLE	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS	RDPS.	FORMATION	RDPS	RADPS
LITHOLOGIC DESCRIPTION									
LITHOLOGIC DESCRIPTION									
FORAN-DIATOR-BEARING CLAYEY NANNO CHALK, light greenish gray (SG 8/1), stiff, severe drilling disturbance, some darker green burrowed segments. <i>Ethmiodiscus rex</i> common.									
SS at CC									
Clay Forams Rads									
CC									
LOWER MIocene									
GLOBALIGERINATA DISSIMILITIS N5									
CYRTOCARPAS TESTICRIPPA									
DISCOASTER DRUGGET N2									
LYCHMECHANOMA ELONGATA									
Site 366 Hole A		Core 24		Cored Interval: 215.0-224.5 m		Core 25		Cored Interval: 224.5-234.0 m	
ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE	ZONES	AGE
LITHOLOGIC DESCRIPTION									
LITHO-SAMPLE									

Site 366 Hole A Core 31 Cored Interval: 281.5-291.0 m		LITHOLOGIC DESCRIPTION	
AGE	ZONES	FOSIL CHARACTER	METERS SECTION
OOLIGOCENE	Sphaeroidellus cf perakensis NP25 Dorcadosyrinx aetuechus	PELAGIC CLAY	0
		INTERBEDDED CLAYEY NANNO CHALK AND PELAGIC CLAY	0.5
		CLAYEY NANO CHALK, light greenish gray (SG 8/1), stiff, slight drilling disturbance, burrowed, Mn spots with fine-grained halos scattered throughout.	1.0
		PELAGIC CLAY, dark greenish gray (SG 6/1), stiff, slight drilling disturbance, burrowed, sometimes laminae.	1.5
		These appear to be dissolution cycles.	2.0
		SSS at CC (dominant lithology)	2.5
		Clay	3.0
		Ran nos	3.5
		Carbonate Bomb: 3-84 to 85 cm = 81%	4.0
			4.5
			5.0

Site 366 Hole A Core 32 Cored Interval: 291.0-300.5 m		LITHOLOGIC DESCRIPTION	
AGE	ZONES	FOSIL CHARACTER	METERS SECTION
OOLIGOCENE	Globigerina cf perakensis P22 Sphaeroidellus cf perakensis NP25 Dorcadosyrinx aetuechus	PELAGIC CLAY	0
		INTERBEDDED CLAYEY NANNO CHALK AND PELAGIC CLAY	0.5
		CLAYEY NANO CHALK, light greenish gray (SG 8/1), stiff, slightly disturbed, Mn spots with fine-grained halos scattered throughout.	1.0
		PELAGIC CLAY, dark greenish gray (SG 6/1), stiff, slight drilling disturbance, burrowed, sometimes laminae.	1.5
		These appear to be dissolution cycles.	2.0
		SSS at CC (dominant lithology)	2.5
		Clay	3.0
		Ran nos	3.5
		Carbonate Bomb: 3-84 to 85 cm = 81%	4.0
			4.5
			5.0

Site 366 Hole A		Core 33	Cored Interval: 300.5-310.0 m
			LITHOLOGIC DESCRIPTION
ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY
AGB		0	VOID
		0.5	
		1	
		1.0	
		2	
		3	
		4	
		5	
		6	
			OOLITHIC NANO CHALK AND PELAGIC CLAY Sphaerodiscus distentus NP21 Dorcadopsispis asteucinus

Site 366 Hole A		Core 34	Cored Interval: 310.0-319.5 m
			LITHOLOGIC DESCRIPTION
ZONES	FOSSIL CHARACTER	METERS	LITHOLOGY
AGB		0	VOID
		0.5	
		1	
		1.0	
		2	
		3	
		4	
		5	
		6	
			OOLITHIC NANO CHALK AND PELAGIC CLAY Sphaerodiscus distentus NP24 Globorotalia optima optima P21 Sphaerodiscus distentus NP24 Dorcadopsispis asteucinus

Site 366		Hole A	Core 36	Cored Interval: 329.0-338.5 m
				LITHO-SAMPLE
				DEFORMATION
ZONES	FOSIL CHARACTER	SECTION	METER	LITHOLOGY
AEF	RADS	0	0.5	VOID
FORAMS	NANNO.		1	1.0
				INTERBEDDED CLAYEY NANNO CHALK AND PELAGIC CLAY
				CLAYEY NANNO CHALK, light greenish gray (5G 8/1), stiff, slight to no drilling disturbance, disturbance, burrowed, Mn bands, laminae, halos abundant, some layers laminated.
				PELAGIC CLAY, dark greenish gray (5G 6/1), stiff, no drilling disturbance, burrowed.
				Sometimes the contact between pelagic clay below clayey nanno is gradational but the other contact is always sharp. These alternations appear to be dissolution cycles.
				SS at CC (dominant lithology) Clay C Forams R Nanno A Fish debris R
				Carbonate Bomb: 3-53 to 54 cm = 69%
				OLOCENE
				Globorotalia optima optima P21 Sphaerolithus distentus NP24 Drepanodiscus reticulatus NP24
				Core Catcher
				CC

Site 366		Hole A	Core 35	Cored Interval: 319.5-329.0 m
				LITHO-SAMPLE
				DEFORMATION
ZONES	FOSIL CHARACTER	SECTION	METER	LITHOLOGY
AEF	RADS	0	0.5	bx
FORAMS	NANNO.	1	1.0	VOID
				INTERBEDDED CLAYEY NANNO CHALK AND PELAGIC CLAY
				CLAYEY NANNO CHALK, light greenish gray (5G 8/1), stiff, slight to no drilling disturbance, disturbance, burrowed, Mn flocs and lithofrag bands scattered throughout, laminated zones present throughout.
				PELAGIC CLAY, dark greenish gray (5G 6/1), stiff, slight to no drilling disturbance, burrowed.
				These alternations appear to be dissolution cycles.
				SS at CC (dominant lithology) Clay C Forams R Nanno A Fish debris R
				Carbonate Bomb: 3-73 to 74 cm = 80%
				OLOCENE
				Globorotalia optima optima P21 Sphaerolithus distentus NP24 Drepanodiscus reticulatus NP24
				Core Catcher
				CC
				N A M-G F A G R -

Site 366 Hole A		Core 38	Cored Interval : 348-0-357.5 m
ZONES	AGE	LITHOLOGY	LITHO. SAMPLE
FARMAS	0	INTERBEDDED CLAYEY MAMMO CHALK AND PELAGIC CLAY	DEFORMATION
MAMMOS	0.5	CLAYEY MAMMO CHALK, light greenish gray (SG 8/1), stiff, no drilling disturbance, burrowed, abundant ln fleck, bands, and lissegang halos, some zones are laminated and are blue-gray and green.	SS at CC (dominant lithology)
RADS	1	PELAGIC CLAY, dark greenish gray (SG 6/1), stiff, no drilling disturbance, burrowed.	Clay A Forams C Nanom A Rads R Fish debris
PRBS.	1.0		These alternations appear to be dissolution cycles.
ABUND.	2		Carbonate Bomb: 3-45 to 46 cm = 85%
FOSIL			
PPBS.			
SCITION			
METERS			
CHARACTER			
FOSSIL			
ABUND.			
PRBS.			
SECTION			
SITE			
0			
1			
2			
3			
4			
			CC
			N C A M
			F A G R F
			Care Catcher
Globorotalia optima P21		Thecocystis tuberosa	
Sphaerotilus distensus NP24		Thecocystis tuberosa	

DEEP SEA DRILLING PROJECT

LEG 41 SITE 367

SITE SUMMARY SHEET

PRINCIPAL RESULTS:

Cape Verde Basin

An 1153-meter section was drilled in the Cape Verde Basin, southeast of the Cape Verde Islands, in the magnetic quiet zone. The hole bottomed in basalt overlain by ?middle to upper Jurassic sediments. It is not entirely clear if the basalt is a sill or if it represents the top of layer 2. Regional correlations and preliminary examination of the rocks, however, support an interpretation suggesting that the basalt represents basement. The Mesozoic part of the section is strikingly similar to the one sampled by Leg 11 in the North American Basin (especially Site 105). This similarity can be observed even in the minute details of sedimentary structures. The differences, minor in importance, are: 1) slightly larger amounts of chert, 2) absence of large scale redeposition features at the Cape Verde Basin site, and 3) the presence of some variegated claystone at the base of the black shale sequence. The entire sedimentary section consists of deep-water deposits. The section was divided into the following units from top to bottom: 1) Pleistocene to Miocene marls and silty clays with occasional sand layers in the Pliocene and other indications of redeposition of originally shallow-water sediment; 2) late Eocene radiolarian clay and middle to early Eocene zeolitic clay and chert; 3) ?Paleocene to Late Cretaceous multicolored silty clay in which the benthonic foraminifera faunas are similar to the Alpine flysch assemblages of the same age; 4) Cenomanian to late Aptian black shales rich in organic matter (organic carbon up to 6.7%) and methane, with abundant pyrite, and rare crystals of partly pyritized barite; 5) Early Aptian to Barremian variegated claystone; 6) Neocomian to Tithonian light gray limestones interbedded with dark gray marls and shales, and rich in aptychi with some ammonites (resembling the sequence outcropping in Maio, Cape Verde Islands); 7) late to ?middle Jurassic reddish brown argillaceous limestones, marls, and claystones with aptychi; and 8) basalt.

Horizon A correlates with middle Eocene cherts. Horizon β possibly correlates with some sideritic beds within the black shale section. The uppermost part of the rather diffuse reflector C correlates with the top of the limestones but no clear correlation can be established between the seismic data and the basalt.

Date occupied	1840Z 3 March 1975
Date departed	0238Z 10 March 1975

LEG 41 SITE 367
SITE SUMMARY SHEET, con't.

Time on site	6 days, 7 hours, 58 minutes
Position: Latitude	12°29.2'N
Longitude	20°02.8'W
Water depth (sea level)	4748 corrected meters
Water depth (rig floor)	4758 corrected meters
Penetration	1153 meters
Number of holes	1
Number of cores	40
Total length of cored section	347.0 meters
Total core recovered	174.3 meters
Percentage core recovery	50
<u>Oldest Sediment Cored</u>	
Depth subbottom	1146 meters
Nature	Red argillaceous limestone
Age	Kimmeridgian to Oxfordian
Measured velocity	~3.5 km/sec
<u>Basement</u>	
Depth subbottom	1146 meters
Nature	Basalt
Velocity range	4.3 km/sec

Site 367 Hole Core 1 Cored Interval : 0-0-8.0 m									
LITHOLOGIC DESCRIPTION									
ZONES	FOSSIL CHARACTER	METER SECTION	LITHOLOGY	PRCS.	ABUND.	FORMS	NANNO	AGE	
0	-	0.5	VOID	R	N				
		1		R	N				
		1.0		R	N				
2		65	10YR 5/2						
3									
4		20							
5		75							
6									

QUATERNARY (unzoned) Geophyrocapsa oceanica N22 - Emiliania huxleyi N22
PLIOTOCENE N22 Geophyrocapsa oceanica N20 - Emiliania huxleyi N21
PLIOTOCENE (unzoned)

Carbonate Bomb: 5-49 to 50 cm = 41%

Site 367 Hole Core 2 Cored interval : 8.0-17.5 m									
LITHOLOGIC DESCRIPTION									
ZONES	FOSSIL CHARACTER	METER SECTION	LITHOLOGY	PRCS.	ABUND.	FORMS	NANNO	AGE	
0	NANNO MARL with various biogenic compositions.	0	VOLO	N					
	NANNO MARL, moderate yellowish brown (10YR 5/2) with mixed grayish brown (2.5Y 5/2), and dark greenish gray (5Gy 4/1) in Section 5, soft, intense drilling disturbance.	1	0.5	N					
	At 4-0 to 40 cm FORAM-BEARING NANNO MARL, medium light gray (N5), soft, intense drilling disturbance.	1	1.0	N					
	At 5-75 to 77 cm SILTY NANNO MARL, dark greenish gray (5Gy 4/1), soft, intense drilling disturbance, Mn stains at upper contact.	2	1.0	N					
	Minor flame feature formed by drilling disturbance at 6.100 to 115 is a FORAM-BEARING NANNO DOME.	2	2	N					
	SS at 1-120 (dominant lithology)	3	2	N					
	Quartz R Diatoms R Heavy minerals R Rads R Forams C Sponge spicules R Fish debris R Nannos A	3	3	N					
	SS at 2-65 (dominant lithology)	4	2	N					
	Quartz C Nannos R Diatoms R Feldspar R Clay A Rads R Nannos A	4	4	N					
	SS at 4-20 (minor lithology)	5	2	N					
	Quartz R Nannos A Diatoms R Forams C Clay A Rads R Nannos A	5	5	N					
	SS at 5-75 (minor lithology)	6	2	N					
	Quartz C Nannos R Diatoms R Heavy minerals R Rads R Clay A Rads R Carbonate unspecificed R Sponge spicules R Nannos R	6	6	N					
	SS at 6-10 (minor lithology)	7	2	N					
	Quartz R Nannos C Rads R Forams C Sponge spicules R Nannos C	7	7	N					
	SS at CC	8	2	N					
	Quartz R Kspar R Heavy minerals C Clay A Glaucite R Rads R Nannos C	8	8	N					
	Carbonate Bomb: 2-63 to 64 cm = 42%	9	2	N					
		9	9	N					
		10	2	N					
		10	10	N					

QUATERNARY (unzoned) Geophyrocapsa oceanica N22 - Emiliania huxleyi N22
PLIOTOCENE N22 Geophyrocapsa oceanica N20 - Emiliania huxleyi N21
PLIOTOCENE (unzoned)

N F P-
F A G Core
R R N Catcher
R R N —

Site 367		Hole	Core 3	Cored Interval: 54.0-53.5 m	Core 4	Cored Interval: 63.5-73.0 m
		LITHOLOGIC DESCRIPTION				
AGE		ZONES FORAMS NANOMS RADOS FOSIL ABUND.	FOSIL CHARACTER	METERS SECTION	LITHOLOGY	LITHOLOGIC DESCRIPTION
				0	N	INTERBEDDED NANO MARL AND CLAYEY SAND.
						NANO MARL, silt, and foram-bearing in some samples, dark yellow brown (10R 4/1) in upper part grading to light brownish gray (2.5Y 6/2) in Section 2, stiff, severe to moderate drilling disturbance, interbeds of FORAM-BEARING NANO CLAYEY SAND, medium grain size.
						Mn stains and flacks concentrated from 2.5M to 6M and 2.8M to 9M.
						Gradational color boundary at 3-7M between overlying and underlying nano marls.
						SS at 1-10 (minor lithology)
						Quartz A
						Diatoms R
						Nanos R
						Rads R
						Forams C
						Clay C
						Forams C
						Quartz A
						Nanos A
						Rads R
						Forams C
						SS at 1-62 (minor lithology)
						Quartz A
						Nanos A
						Diatoms A
						Clay A
						Forams C
						Quartz A
						Nanos A
						Rads R
						Forams C
						SS at 2-30 (dominant lithology)
						Quartz A
						Nanos A
						Rads R
						Forams C
						Quartz A
						Nanos A
						Rads R
						Forams C
						SS at 3-28 (dominant lithology)
						Quartz A
						Nanos A
						Rads R
						Forams A
						Clay A
						Calcareous R
						Rads R
						Forams C
						Quartz A
						Nanos A
						Rads R
						Forams C
						SS at 3-90 (minor lithology)
						Quartz A
						Nanos A
						Rads R
						Forams C
						Clay A
						Calcareous R
						Rads R
						Forams C
						Quartz A
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A
						Calcite rhombs A
						Rads R
						Forams R
						Quartz R
						Nanos A
						Rads R
						Forams C
						SS at 4-40 (dominant lithology)
						Quartz R
						Forams R
						Heavy minerals R
						Nanos A
						Clay A</

Site 367		Hole	Core 5	Cored Interval : 150.5-160.0 m
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N F-C M	0		QUARTZ SAND, pale yellowish brown (10R 6/2), firm, intense drilling disturbance.	
F F G	0.5		Marl, grayish olive (10Y 4/2), firm, intense drilling disturbance, Mn stains.	
R R -	1.0		SS at 1-82 (dominant lithology)	
			Heavy minerals R	R
			Forams D	Glaucite
			Forams R	R
			Rads A	R
			Clay A	R
			NaNos A	R
			Quartz R	R
* Ceratolithus rugosus MN13				

Site 367		Hole	Core 6	Cored Interval : 236.0-245.5 m
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at 1-76 (dominant lithology)	
			Quartz C	D
			Heavy minerals R	R
			Clay R	R
			Fish debris R	R
			SS at 1-117 (dominant lithology)	
			Quartz C	D
			Heavy minerals R	A
			Glaucite R	R
			CC	
MIDDLE Miocene				

Site 367		Hole	Core 7	Cored Interval : 245.5-255.0 m
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R C M	0		SILT-BEARING CLAY, dark greenish gray (5G 4/1), stiff, greasy, severe drilling disturbance.	
F F G	0.5		VOID	
R R -	1.0		1	1.0-
			1	0.5-
			0	0.5-
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOLITE, light olive gray (5Y 5/2).	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Glaucite R	
			NaNos C	
			Clay R	
LITHOLOGY				
DEFORMATION				
LITHO. SAMPLE				
AGE				
Site 367				
ZONES	AGES	FOSSIL FORMAMS	LITHOLOGY	METERS SECTION
N R P	0		VOID	
F A M	0.5		NaNos N	
R R -	1.0		SILSTONE, olive gray (5Y 3/2) and a FERRODOL	

71

Site 367		Core 22		Cored Interval: 720.5-730.0 m	
Hole	Zones	Fossil Character	Fossil	Section	Lithology
AGE		PRES.	PRES.	METERS	DEFORMATION
					LITHO. ASMPLE
					LITHOLOGY
	0				ALTERNATIONS OF SHALE, CLAY, AND SILT TURBIDITES.
					NANNO-BEARING CLAY, dark gray black (N3) to dark greenish black (5G 2/1) to dark greenish gray (5G 4/1), hard, burned, faint laminations, severe drilling disturbance (broken).
					SHALE, black (N1), fissile, gas-bearing, burned, severe drilling disturbance.
					Clays are interbedded with thin quartz silt turbidites (1 cm to 12 cm thick) scattered throughout the core.
					SS at 2-37 (dominant lithology)
					Quartz R
					Clay R
					Nannos R
					Fe/Mn nodules R
					Rads C
					SS at 2-97 (dominant lithology)
					Heavy minerals R
					Clay A
					Carbonate R
					Forams R
					Crystals C
					Rads C
					Pyrite C
					Fe/Mn nodules C
					SS at 2-138 (minor lithology)
					Quartz R
					Clay R
					Fe/Mn par R
					Fish debris R
					Heavy minerals R
					SS at 3-51 (minor lithology)
					Dolomite R
					Clay R
					Nannos R
					SS at 5-78 (minor lithology)
					Quartz C
					Clay A
					Nannos C
					Pyrite C
					Cone Catcher C
	0				78
	1				6
	2				5
	3				4
	4				3
	5				2
	6				1
	7				0
					37
					97

Site 367 Hole		Core 27		Cored Interval : 939.0-948.5 m	
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION	
	0		VOID	ALTERATIONS OF LIGHT GRAY LIMESTONE AND OLIVE BLACK TO OLIVE GRAY SHALE. SHALE, olive black (5Y 2/1), fissile, gas-bearing, disseminated pyrite, a few apytychi found.	
	1	0.5		NANNO LIMESTONE, 1 light gray (N7), burrowed, Mn streaks common, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common, Mn streaks common.
	2	1.0		NANNO LIMESTONE, 1 varicolored olive gray (5Y 4/1), finely laminated, gas-bearing, apytychi-bearing.	NANNO LIMESTONE, 1 varicolored olive gray (5Y 4/1), finely laminated, gas-bearing, apytychi-bearing.
	3	C P-M		All contacts are fairly sharp although some show diffusion? of colors into the surrounding limestone.	All contacts are fairly sharp although some show diffusion? of colors into the surrounding limestone.
	F —	R R	Core Catcher	SS at 1-131 (minor lithology) Quartz R Heavy minerals A Clay R SS at 3-94 (minor lithology) Pyrite A Fe/Mn C Nannos C Carbone Bomb: 3-84 to 85 cm = 92%	SS at 1-131 (minor lithology) Quartz R Glaucite N Heavy minerals A Nannos A Clay A Fe/Mn R SS at 3-60 (dominant lithology) Calcareous crystals A Forams Nannos R Carbone Bomb: 3-84 to 85 cm = 92%

Site 367 Hole		Core 30		Cored Interval : 1024.5-1034.0 m	
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION	
	0			NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	1	0.5		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	2	1.0		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	3	C P-M		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	F —	R R	Core Catcher	SS at 1-10 (minor lithology) Quartz R Albite A Unspecified A Nannos C Carbone Bomb: 1-30 to 101 cm = 96%	SS at 1-10 (minor lithology) Quartz R Albite A Unspecified A Nannos C Carbone Bomb: 1-30 to 101 cm = 96%

Site 367 Hole		Core 29		Cored Interval : 996.0-1005.5 m	
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION	
	0		VOID	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common, occasional CHELT modules, medium gray (N5).	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common, occasional CHELT modules, medium gray (N5).
	1	0.5		Thin bed of NANNO MARLSTONE, varicolored olive gray (5Y 4/1), finely laminated. Occurs at 2-38 to 43 cm.	Thin bed of NANNO MARLSTONE, varicolored olive gray (5Y 4/1), finely laminated. Occurs at 2-38 to 43 cm.
	2	1.0		SS at 1-55 (minor lithology) Quartz C Calcite A Nannos C	SS at 1-55 (minor lithology) Quartz C Calcite A Nannos C
	3	C P-M		Carbone Bomb: 2-40 to 41 cm = 92% 2-106 to 107 cm = 96%	Carbone Bomb: 2-40 to 41 cm = 92% 2-106 to 107 cm = 96%
	F —	R R	Core Catcher		

Site 367 Hole		Core 28		Cored Interval : 967.5-977.0 m	
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION	
	0		VOID	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	1	0.5		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	2	1.0		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	3	C P-M		SS at 2-22 (minor lithology) Pyrite R Clay R Nannos R Carbone Bomb: 2-48 cm = 96%	SS at 2-22 (minor lithology) Pyrite R Clay R Nannos R Carbone Bomb: 2-48 cm = 96%
	F —	R R	Core Catcher	SS at 2-79 (dominant lithology) Nannos C	SS at 2-79 (dominant lithology) Nannos C

Site 367 Hole		Core 27		Cored Interval : 939.0-948.5 m	
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION	
	0		VOID	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	1	0.5		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	2	1.0		NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.	NANNO-BEARING LIMESTONE, light gray (N7), burrowed, Mn streaks common, Mn streaks common.
	3	C P-M		SS at 2-133 (minor lithology) Quartz R Carbonate C Unspecified D	SS at 2-133 (minor lithology) Quartz R Carbonate C Unspecified D
	F —	R R	Core Catcher	SS at CC Carbonate R Unspecified D	SS at CC Carbonate R Unspecified D

Site 367		Hole	Core 38	Cored Interval : 1142.0-1148.0 m
ZONES	FOSSIL CHARACTER			
AGE	UPPER JURASSIC			
NOT YOUNGER THAN KIMMERIDGIAN	Fossils RADS NANNO FORAMS			LITHOLOGIC DESCRIPTION
				DEFORMATION
				LITHO. SAMPLE
				LITHOLOGY
			METERS	
			SECTION	
			PRES.	
			ABUND.	
			RADS	
			NANNO	
			FORAMS	
			AGE	

N R-F P

0 VOID

0.5

1.0

1.0 138

1.0 35

2

3

Core Catcher

SS at 1-138 (dominant lithology)
Clay C
Carbonate R
unspecified A
Nanmos C

SS at 2-35 (dominant lithology)
Chlorite R
Carbonate Clay
unspecified A
Nanmos R

Possible flow units: 2-27 to 3-33 cm
3-55 to 3-156 cm
3-105 to 3-150 cm

Site 367		Hole	Core 40	Cored Interval : 1151.0-1153.0 m
ZONES	FOSSIL CHARACTER			
AGE	UPPER JURASSIC			
			LITHOLOGIC DESCRIPTION	
			DEFORMATION	
			LITHO. SAMPLE	
			LITHOLOGY	
			METERS	
			SECTION	
			PRES.	
			ABUND.	
			RADS	
			NANNO	
			FORAMS	
			AGE	

0 0.5

1 1.0

Core Catcher

BASALT, black (N), calcite veined and chlorite, chert clast at 40 cm, disseminated pyrite throughout, amygdaloidal with green mineral and pyrite and glass fillings. Grain size larger than in Core 40.

Site 367		Hole	Core 39	Cored Interval : 1148.0-1151.0 m
ZONES	FOSSIL CHARACTER			
AGE	UPPER JURASSIC			
			LITHOLOGIC DESCRIPTION	
			DEFORMATION	
			LITHO. SAMPLE	
			LITHOLOGY	
			METERS	
			SECTION	
			PRES.	
			ABUND.	
			RADS	
			NANNO	
			FORAMS	
			AGE	

0 VOID

0.5

1 1.0

Core Catcher

BASALT, breccia clasts of unaltered, thick chlorite zones, calcite veins, amygdaloids filled with clear crystals and/or white, soft material, common pyrite disseminated throughout, chert clast at 118 cm.

DEEP SEA DRILLING PROJECT

LEG 41 SITE 368

SITE SUMMARY SHEET

PRINCIPAL RESULTS:

Cape Verde Rise

A sedimentary section of Cape Verde Rise was sampled down to 984.5 meters, the level of a sharp deep reflector. The sediments are predominantly terrigenous and almost barren of microfossils, except in the upper part and near the base of the section. The deep reflector was found to correspond with diabase sills interstratified within black shales near the top of that sedimentary unit. The sediment section consists of the following units (from top to bottom): 1) Quaternary to middle Miocene nannofossil oozes and marls with volcanic ash layers in the upper Miocene; 2) early Miocene to ?early Eocene green clays and claystones with chert and porcellanite; 3) ?Paleocene to ?late Cretaceous red and green shales; 4) ?late Cretaceous green claystone and shales; and 5) ?Turonian to Albian black shales with diabase sills. Three observations strongly suggest that the Cape Verde Rise resulted from a broad uplift of about 1000 to 1500 meters that could have taken place in the late Paleogene to early Neogene: 1) the occurrence of basic sills, probably related to Miocene volcanic activity on Cape Verde Islands; 2) abundant turbidite deposition in the Paleogene and late Cretaceous; and 3) the occurrence of carbonates restricted to the upper part of the section.

Date occupied	0651 hrs March 13, 1975
Date departed	1928 hrs March 20, 1975
Time on site	7 days, 2 hours, 37 minutes
Position: Latitude	17°30.4'N
Longitude	21°21.2'W
Water depth (sea level)	3366 corrected meters
Water depth (rig floor)	4758 corrected meters
Penetration	984.5 meters
Number of holes	1
Number of cores	63

LEG 41 SITE 368
SITE SUMMARY SHEET, con't.

Total length of cored section 582.5 meters

Total core recovered 327.7 meters

Percentage core recovery 56%

Oldest Sediment Cored

Depth subbottom 984.5 meters

Nature Black shale



Age Albian

Measured Velocity 3.06 km/sec

83

Site 368		Hole	Core 3	Cored Interval: 47.0-56.5 m
				LITHOLOGIC DESCRIPTION
ZONES	FOSIL CHARACTER	METRES	SECTION	LITHOLOGY
FORAMS		0	N	FORAM MUD MARL AND Ooze, yellow gray (SY 8/1), gray green (SY 5/1), soft, intense drilling disturbance, mottled with black and greenish gray (SY 6/1).
NANNO.		0.5		SS at 1-98 (dominant lithology)
RADS		1		Quartz R
ABUND.		1.0		Clay A
FOSIL		1.0		Heavy minerals R
NANNO.		1.0		Forams C
AGE		1.0		Nannos A
SS at 2-80 (dominant lithology)				
Clay	C	80		Forams A
Nanmos	A			Diatoms R
Mollusk	R			
SS at 4-120 (minor lithology)				
Mic	R			Heavy minerals R
Clay	A			Fe/Mn R
Forams	A			Nanmos A
Mollusk	R			Phosphate R
SS at 6-102 (dominant lithology)				
Heavy minerals	R			Clay A
Forams	C			Nanmos A
Rads	R			Sponge spicules R
Plant debris	R			
Carbonate Bomb: 3-80 to 81 cm = 68%				
Carbonate Bomb: 3-82 to 83 cm = 65%				
LOWER PLEISTOCENE				
Globorotalia truncatuloides Zone (Globorotalia crassiformis violacea Subzone)				
Pseudodemnum titanum Lacunoosa NNT9				
Globorotalia marginata Margaritace Zone (Globorotalia offenerstera pseudomulticostata NNT5)				
Discaster asymmetricus NNT4-Reticulofenestrata pseudomulticostata NNT5				
- - -				
R	P			
N	A			
F	G			
N	A			
Core Catcher				
CC				

Site 368		Hole	Core 4	Cored Interval: 85.0-94.5 m
				LITHOLOGIC DESCRIPTION
ZONES	FOSIL CHARACTER	METRES	SECTION	LITHOLOGY
FORAMS		0	N	FORAM-BEARING MAMMO MARL AND Ooze, very light gray (NB), light gray (SY 6/1), soft, to severe drilling disturbance, streaks of black throughout, mottled, a cemented burrow occurs at 4-71 cm.
NANNO.		0.5		SS at 1-80 (dominant lithology)
RADS		1		Quartz R
ABUND.		1.0		Clay A
FOSIL		1.0		Heavy minerals R
NANNO.		1.0		Forams C
AGE		1.0		Nannos A
SS at 2-65 (dominant lithology)				
Clay	R	80		Carbonate R
Nanmos	A			Unspecified R
Mollusk	R			Sponge spicules R
SS at 3-65 (minor lithology)				
Mic	R			Quartz Fe/Mn R
Clay	A			Nanmos R
Forams	A			Forams R
Mollusk	R			Rads R
SS at 5-70 (dominant lithology)				
Clay	C	65		Clay C
Nanmos	A			Nanmos A
Rads	R			Phosphate R
SS at 6-55 (dominant lithology)				
Quartz	R			Quartz R
Mica	R			Mica R
Heavy minerals	R			Heavy minerals R
Forams	C			Forams C
Rads	R			Rads A
Carbonate Bomb: 3-82 to 83 cm = 65%				
LOWER PLIOCENE				
Globorotalia marginata Margaritace Zone (Globorotalia offenerstera pseudomulticostata NNT5)				
- - -				
R	P			
N	A			
F	G			
N	A			
Core Catcher				
CC				



Site 368		Hole	Core 9	Cored Interval: 189.5-199.0 m
				LITHOLOGIC DESCRIPTION
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	
METERS	RADS	PRES.		
ABUND.	NANNO	FOSIL		
FORMS	NA	CHAR		
0			INTERBEDDED NANNO MARL AND NANNO-BEARING CLAY	
			NANNO MARL, yellow gray (SY 8/1), stiff, slight drilling disturbance, bioturbated, black specks scattered throughout. Gradational color boundaries, because of bioturbation, with	
			NANNO-BEARING CLAY, various shades of yellow brown (10R 5/4), stiff, bioturbated.	
			SS at 2-43 (dominant lithology)	
			Quartz R	
			Clay A	
			Forams R	
			Carbonate A	
			Unspecified R	
			Nanomos A	
			SS at 3-90 (dominant lithology)	
			Quartz R	
			Heavy minerals R	
			Forams C	
			Rads R	
			Nanomos A	
			SS at 5-42 (dominant lithology)	
			Quartz R	
			Heavy minerals R	
			Nanomos C	
			SS at CC	
			Clay R	
			Fish debris R	
			SS at CC	
			Quartz R	
			Heavy minerals R	
			Forams R	
			Carbonate Bomb: 3-7 to 8 cm = 25%	

Site 368		Hole	Core 10	Cored Interval: 199.0-208.5 m
				LITHOLOGIC DESCRIPTION
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	
METERS	RADS	PRES.		
ABUND.	NANNO	FOSIL		
FORMS	NA	CHAR		
0			INTERBEDDED NANNO MARL AND NANNO-BEARING CLAY	
			NANNO MARL, light greenish gray (SY 8/1), stiff, slight to no drilling disturbance, bioturbated.	
			NANNO-BEARING CLAY, moderate brown (SYR 4/4), stiff, slight to no drilling disturbance, bioturbated.	
			Color banding at intervals of 5 to 15 cm.	
			SS at 1-60 (dominant lithology)	
			Quartz C	
			Mica R	
			Heavy minerals R	
			Clay A	
			Pyrite R	
			Fe/Mn R	
			Nanomos C	
			SS at 1-70 (minor lithology)	
			Quartz R	
			Feldspar R	
			Clay A	
			Pyrite R	
			Forams R	
			Rads R	
			Molusks R	
			SS at CC	
			Quartz C	
			Mica R	
			Clay A	
			Hematite R	
			Nanomos R	
			Carbonate Bomb: 2-52 to 53 cm = 13%	

Site 368		Hole	Core 11	Cored Interval: 208.5-218.0 m
				LITHOLOGIC DESCRIPTION
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	
METERS	RADS	PRES.		
ABUND.	NANNO	FOSIL		
FORMS	NA	CHAR		
0			INTERBEDDED SILTY CLAY AND NANNO Ooze	
			SILTY CLAY, wide variation of greens, grays, blues, and browns, dominant colors are light and medium blue gray (SB 7/1 to SB 5/1); firm, slight to intense drilling disturbance, bioturbated.	
			NANNO Ooze, light bluish gray (SB 7/1), firm, slight drilling disturbance, bioturbated.	
			Color breaks between lithologies are sharp to gradational.	
			SS at 2-68 (dominant lithology)	
			Quartz C	
			Mica R	
			Heavy minerals R	
			Clay A	
			Pyrite R	
			Carbonate R	
			unspecified R	
			SS at 2-103 (minor lithology)	
			Heavy minerals R	
			Clay C	
			Nanomos A	
			SS at 3-38 (minor lithology)	
			Quartz C	
			Mica R	
			Feldspar A	
			Clay A	
			Pyrite R	
			Carbone R	
			Mn C	
			Rads C	
			Forams R	
			Nanomos R	
			Carbonate Bomb: 3-52 to 53 cm = 31%	

Core 16		Core Interval: 256.0-265.5 m			
Site 368	Hole	AGE	ZONES	FOSIL	LITHOLOGY
		METERS	SECTIONS	PRBS.	DETERM.
				RADS.	LITHO. SAMPLE
		0	0	VOID	NANNO MARL WITH MINOR INTERBEDS OF CLAY
		0.5	1	N	NANNO MARL; medium bluish gray (SB 5/1), calcareous; stiff, moderate coring disturbance. Section 4 is color banded with interbeds of
		1.0	1	N	C.L.A.; green (SG 6/1), stiff, moderate drilling disturbance. Occurs at 4.5 to 10, 4.15 to 22, 4.42 to 49, 4.88 to 96, and 4.144 to 150.
		56.6/1	2	N	SS at 3-120 (dominant lithology) Quartz R Clay R Carbonate A unspecified R Sponge spicules R
		56.6/1	3	bx	SS at 4-145 (minor lithology) Quartz R Clay D Carbonate R unspecified R Sponge spicules R
		56.6/1	4	bx	bx
		56.6/1	5	bx	bx
		56.6/1	6	bx	bx
		56.6/1	7	bx	bx
		56.6/1	8	bx	bx
		56.6/1	9	bx	bx
		56.6/1	10	bx	bx
		56.6/1	11	bx	bx
		56.6/1	12	bx	bx
		56.6/1	13	bx	bx
		56.6/1	14	bx	bx
		56.6/1	15	bx	bx
		56.6/1	16	bx	bx
		56.6/1	17	bx	bx
		56.6/1	18	bx	bx
		56.6/1	19	bx	bx
		56.6/1	20	bx	bx
		56.6/1	21	bx	bx
		56.6/1	22	bx	bx
		56.6/1	23	bx	bx
		56.6/1	24	bx	bx
		56.6/1	25	bx	bx
		56.6/1	26	bx	bx
		56.6/1	27	bx	bx
		56.6/1	28	bx	bx
		56.6/1	29	bx	bx
		56.6/1	30	bx	bx
		56.6/1	31	bx	bx
		56.6/1	32	bx	bx
		56.6/1	33	bx	bx
		56.6/1	34	bx	bx
		56.6/1	35	bx	bx
		56.6/1	36	bx	bx
		56.6/1	37	bx	bx
		56.6/1	38	bx	bx
		56.6/1	39	bx	bx
		56.6/1	40	bx	bx
		56.6/1	41	bx	bx
		56.6/1	42	bx	bx
		56.6/1	43	bx	bx
		56.6/1	44	bx	bx
		56.6/1	45	bx	bx
		56.6/1	46	bx	bx
		56.6/1	47	bx	bx
		56.6/1	48	bx	bx
		56.6/1	49	bx	bx
		56.6/1	50	bx	bx
		56.6/1	51	bx	bx
		56.6/1	52	bx	bx
		56.6/1	53	bx	bx
		56.6/1	54	bx	bx
		56.6/1	55	bx	bx
		56.6/1	56	bx	bx
		56.6/1	57	bx	bx
		56.6/1	58	bx	bx
		56.6/1	59	bx	bx
		56.6/1	60	bx	bx
		56.6/1	61	bx	bx
		56.6/1	62	bx	bx
		56.6/1	63	bx	bx
		56.6/1	64	bx	bx
		56.6/1	65	bx	bx
		56.6/1	66	bx	bx
		56.6/1	67	bx	bx
		56.6/1	68	bx	bx
		56.6/1	69	bx	bx
		56.6/1	70	bx	bx
		56.6/1	71	bx	bx
		56.6/1	72	bx	bx
		56.6/1	73	bx	bx
		56.6/1	74	bx	bx
		56.6/1	75	bx	bx
		56.6/1	76	bx	bx
		56.6/1	77	bx	bx
		56.6/1	78	bx	bx
		56.6/1	79	bx	bx
		56.6/1	80	bx	bx
		56.6/1	81	bx	bx
		56.6/1	82	bx	bx
		56.6/1	83	bx	bx
		56.6/1	84	bx	bx
		56.6/1	85	bx	bx
		56.6/1	86	bx	bx
		56.6/1	87	bx	bx
		56.6/1	88	bx	bx
		56.6/1	89	bx	bx
		56.6/1	90	bx	bx
		56.6/1	91	bx	bx
		56.6/1	92	bx	bx
		56.6/1	93	bx	bx
		56.6/1	94	bx	bx
		56.6/1	95	bx	bx
		56.6/1	96	bx	bx
		56.6/1	97	bx	bx
		56.6/1	98	bx	bx
		56.6/1	99	bx	bx
		56.6/1	100	bx	bx
		56.6/1	101	bx	bx
		56.6/1	102	bx	bx
		56.6/1	103	bx	bx
		56.6/1	104	bx	bx
		56.6/1	105	bx	bx
		56.6/1	106	bx	bx
		56.6/1	107	bx	bx
		56.6/1	108	bx	bx
		56.6/1	109	bx	bx
		56.6/1	110	bx	bx
		56.6/1	111	bx	bx
		56.6/1	112	bx	bx
		56.6/1	113	bx	bx
		56.6/1	114	bx	bx
		56.6/1	115	bx	bx
		56.6/1	116	bx	bx
		56.6/1	117	bx	bx
		56.6/1	118	bx	bx
		56.6/1	119	bx	bx
		56.6/1	120	bx	bx
		56.6/1	121	bx	bx
		56.6/1	122	bx	bx
		56.6/1	123	bx	bx
		56.6/1	124	bx	bx
		56.6/1	125	bx	bx
		56.6/1	126	bx	bx
		56.6/1	127	bx	bx
		56.6/1	128	bx	bx
		56.6/1	129	bx	bx
		56.6/1	130	bx	bx
		56.6/1	131	bx	bx
		56.6/1	132	bx	bx
		56.6/1	133	bx	bx
		56.6/1	134	bx	bx
		56.6/1	135	bx	bx
		56.6/1	136	bx	bx
		56.6/1	137	bx	bx
		56.6/1	138	bx	bx
		56.6/1	139	bx	bx
		56.6/1	140	bx	bx
		56.6/1	141	bx	bx
		56.6/1	142	bx	bx
		56.6/1	143	bx	bx
		56.6/1	144	bx	bx
		56.6/1	145	bx	bx
		56.6/1	146	bx	bx
		56.6/1	147	bx	bx
		56.6/1	148	bx	bx
		56.6/1	149	bx	bx
		56.6/1	150	bx	bx
		56.6/1	151	bx	bx
		56.6/1	152	bx	bx
		56.6/1	153	bx	bx
		56.6/1	154	bx	bx
		56.6/1	155	bx	bx
		56.6/1	156	bx	bx
		56.6/1	157	bx	bx
		56.6/1	158	bx	bx
		56.6/1	159	bx	bx
		56.6/1	160	bx	bx
		56.6/1	161	bx	bx
		56.6/1	162	bx	bx
		56.6/1	163	bx	bx
		56.6/1	164	bx	bx
		56.6/1	165	bx	bx
		56.6/1	166	bx	bx
		56.6/1	167	bx	bx
		56.6/1	168	bx	bx
		56.6/1	169	bx	bx
		56.6/1	170	bx	bx
		56.6/1	171	bx	bx
		56.6/1	172	bx	bx
		56.6/1	173	bx	bx
		56.6/1	174	bx	bx
		56.6/1	175	bx	bx
		56.6/1	176	bx	bx
		56.6/1	177	bx	bx
		56.6/1	178	bx	bx
		56.6/1	179	bx	bx
		56.6/1	180	bx	bx
		56.6/1	181	bx	bx
		56.6/1	182	bx	bx
		56.6/1	183	bx	bx
		56.6/1	184	bx	bx
		56.6/1	185	bx	bx
		56.6/1	186	bx	bx
		56.6/1	187	bx	bx
		56.6/1	188	bx	bx
		56.6/1	189	bx	bx
		56.6/1	190	bx	bx
		56.6/1	191	bx	bx
		56.6/1	192	bx	bx
		56.6/1	193	bx	bx
		56.6/1	194	bx	bx
		56.6/1	195	bx	bx
		56.6/1	196	bx	bx
		56.6/1	197	bx	bx
		56.6/1	198	bx	bx
		56.6/1	199	bx	bx
		56.6/1	200	bx	bx
		56.6/1	201	bx	bx
		56.6/1	202	bx	bx
		56.6/1	203	bx	bx
		56.6/1	204	bx	bx
		56.6/1	205	bx	bx
		56.6/1	206	bx	bx
		56.6/1	207	bx	bx
		56.6/1	208	bx	bx
		56.6/1	209	bx	bx
		56.6/1	210	bx	bx
		56.6/1	211	bx	bx
		56.6/1	212	bx	bx
		56.6/1	213	bx	bx
		56.6/1	214	bx	bx
		56.6/1	215	bx	bx
		56.6/1	216	bx	bx
		56.6/1	217	bx	bx
		56.6/1	218	bx	bx
		56.6/1	219	bx	bx
		56.6/1	220	bx	bx
		56.6/1	221	bx	bx
		56.6/1	222	bx	bx
		56.6/1	223	bx	bx
		56.6/1	224	bx	bx
		56.6/1	225	bx	bx
		56.6/1	226	bx	bx
		56.6/1	227	bx	bx
		56.6/1	228	bx	bx
		56.6/1	229	bx	bx
		56.6/1	230	bx	bx
		56.6/1	231	bx	bx
		56.6/1	232	bx	bx
		56.6/1	233	bx	bx
		56.6/1	234	bx	bx
		56.6/1	235	bx	bx
		56.6/1	236	bx	bx
		56.6/1	237	bx	bx
		56.6/1	238	bx	bx
		56.6/1	239	bx	bx
		56.6/1	240	bx	bx
		56.6/1	241	bx	bx
		56.6/1	242	bx	bx
		56.6/1	243	bx	bx
		56.6/1	244	bx	bx
		56.6/1	245	bx	bx
		56.6/1	246	bx	bx
		56.6/1	247	bx	bx
		56.6/1	248	bx	bx
		56.6/1	249	bx	bx
		56.6/1	250	bx	bx
		56.6/1	251	bx	bx
		56.6/1	252	bx	bx
		56.6/1	253	bx	bx
		56.6/1	254	bx	bx
		56.6/1	255	bx	bx
		56.6/1	256	bx	bx
		56.6/1	257	bx	bx
		56.6/1	258	bx	bx
		56.6/1	259	bx	bx
		56.6/1	260	bx	bx
		56.6/1	261	bx	bx
		56.6/1	262	bx	bx
		56.6/1	263	bx	bx
		56.6/1	264	bx	bx
		56.6/1	265	bx	bx
		56.6/1	266	bx	bx
		56.6/1	267	bx	bx
		56.6/1	268	bx	bx
		56.6/1	269	bx	bx
		56.6/1	270	bx	bx
		56.6/1	271	bx	bx

90

ai

Site 368		Core 20	Cored Interval : 294.0-303.5 m		
A&E	ZONES	FOSIL CHARACTER	FOSILS	METRES	LITHOLOGY
MIDDLE EOCENE? - LOWER MIOCENE?	ADMAMS FORMS	RDS	RESD.	0	SILT CLAY, blue green (BG 5/2), very stiff, moderate drilling disturbance, burrowed.
-	N	R	VOID	0.5	
-	-	F		1	
-	-			1.0	
					Core Catcher

Site 368		Core 22		Cored Interval: 322.5-332.0 m			
Age	Fossils	Section	Metres	Lithology	Deformation	Litho. Sample	Lithologic Description
		0	0.5				EACH SECTION OF THIS CORE EXPANDED BEYOND THE 150 cm CUT. - SOME EXPANDED AS MUCH AS 10 cm.
		1	1.0				CLAYSTONE; medium bluish gray (SB 5/1), grading with depth to thinly interbedded green gray (SG 6/1 and SG 4/1), hard, interbedded with good zoophycos and clionid traces, lessening bands of Mn (?) at some burrows, joint faintly laminated, fracturing caused by core expansion.
		2					In Sections 1 and 2 the medium blue gray color predominates. Beginning with Section 3 the gray (SG 4/1) color predominates with thin (<5 cm) interbeds of SG 6/1. Contacts between colors are sharp.
		3					SS at 1-17 (minor lithology)
		4					Quartz R Clay R Dolomite rhombs D Volcanic glass R
		5					SS at 1-46 (minor lithology)
		6					Quartz R Parite R Dolomite R Fe/Mn Fish debris R
		7					SS at 1-71 (minor lithology)
		8					Quartz D Heavy minerals R Feldspar C Clay C
		9					SS at 2-30 (dominant lithology)
		10					Quartz D Volcanic glass R Pyrite R Carbonate Fish debris R
		11					SS at 2-83 (dominant lithology)
		12					Quartz D Nica R Clay C Carbonate Fish debris R
		13					SS at 2-143 (dominant lithology)
		14					Quartz R Volcanic glass R Pyrite R Fe/Mn R Nanios R Fish debris R
		15					Unspecified R
		16					Unspecified R
		17					Unspecified R
		18					Unspecified R
		19					Unspecified R
		20					Unspecified R
		21					Unspecified R
		22					Unspecified R
		23					Unspecified R
		24					Unspecified R
		25					Unspecified R
		26					Unspecified R
		27					Unspecified R
		28					Unspecified R
		29					Unspecified R
		30					Unspecified R
		31					Unspecified R
		32					Unspecified R
		33					Unspecified R
		34					Unspecified R
		35					Unspecified R
		36					Unspecified R
		37					Unspecified R
		38					Unspecified R
		39					Unspecified R
		40					Unspecified R
		41					Unspecified R
		42					Unspecified R
		43					Unspecified R
		44					Unspecified R
		45					Unspecified R
		46					Unspecified R
		47					Unspecified R
		48					Unspecified R
		49					Unspecified R
		50					Unspecified R
		51					Unspecified R
		52					Unspecified R
		53					Unspecified R
		54					Unspecified R
		55					Unspecified R
		56					Unspecified R
		57					Unspecified R
		58					Unspecified R
		59					Unspecified R
		60					Unspecified R
		61					Unspecified R
		62					Unspecified R
		63					Unspecified R
		64					Unspecified R
		65					Unspecified R
		66					Unspecified R
		67					Unspecified R
		68					Unspecified R
		69					Unspecified R
		70					Unspecified R
		71					Unspecified R
		72					Unspecified R
		73					Unspecified R
		74					Unspecified R
		75					Unspecified R
		76					Unspecified R
		77					Unspecified R
		78					Unspecified R
		79					Unspecified R
		80					Unspecified R
		81					Unspecified R
		82					Unspecified R
		83					Unspecified R
		84					Unspecified R
		85					Unspecified R
		86					Unspecified R
		87					Unspecified R
		88					Unspecified R
		89					Unspecified R
		90					Unspecified R
		91					Unspecified R
		92					Unspecified R
		93					Unspecified R
		94					Unspecified R
		95					Unspecified R
		96					Unspecified R
		97					Unspecified R
		98					Unspecified R
		99					Unspecified R
		100					Unspecified R
		101					Unspecified R
		102					Unspecified R
		103					Unspecified R
		104					Unspecified R
		105					Unspecified R
		106					Unspecified R
		107					Unspecified R
		108					Unspecified R
		109					Unspecified R
		110					Unspecified R
		111					Unspecified R
		112					Unspecified R
		113					Unspecified R
		114					Unspecified R
		115					Unspecified R
		116					Unspecified R
		117					Unspecified R
		118					Unspecified R
		119					Unspecified R
		120					Unspecified R
		121					Unspecified R
		122					Unspecified R
		123					Unspecified R
		124					Unspecified R
		125					Unspecified R
		126					Unspecified R
		127					Unspecified R
		128					Unspecified R
		129					Unspecified R
		130					Unspecified R
		131					Unspecified R
		132					Unspecified R
		133					Unspecified R
		134					Unspecified R
		135					Unspecified R
		136					Unspecified R
		137					Unspecified R
		138					Unspecified R
		139					Unspecified R
		140					Unspecified R
		141					Unspecified R
		142					Unspecified R
		143					Unspecified R
		144					Unspecified R
		145					Unspecified R
		146					Unspecified R
		147					Unspecified R
		148					Unspecified R
		149					Unspecified R
		150					Unspecified R
		151					Unspecified R
		152					Unspecified R
		153					Unspecified R
		154					Unspecified R
		155					Unspecified R
		156					Unspecified R
		157					Unspecified R
		158					Unspecified R
		159					Unspecified R
		160					Unspecified R
		161					Unspecified R
		162					Unspecified R
		163					Unspecified R
		164					Unspecified R
		165					Unspecified R
		166					Unspecified R
		167					Unspecified R
		168					Unspecified R
		169					Unspecified R
		170					Unspecified R
		171					Unspecified R
		172					Unspecified R
		173					Unspecified R
		174					Unspecified R
		175					Unspecified R
		176					Unspecified R
		177					Unspecified R
		178					Unspecified R
		179					Unspecified R
		180					Unspecified R
		181					Unspecified R
		182					Unspecified R
		183					Unspecified R
		184					Unspecified R
		185					Unspecified R
		186					Unspecified R
		187					Unspecified R
		188					Unspecified R
		189					Unspecified R
		190					Unspecified R
		191					Unspecified R
		192					Unspecified R
		193					Unspecified R
		194					Unspecified R
		195					Unspecified R
		196					Unspecified R
		197					Unspecified R
		198					Unspecified R
		199					Unspecified R
		200					Unspecified R
		201					Unspecified R
		202					Unspecified R
		203					Unspecified R
		204					Unspecified R
		205					Unspecified R
		206					Unspecified R
		207					Unspecified R
		208					Unspecified R
		209					Unspecified R
		210					Unspecified R
		211					Unspecified R
		212					Unspecified R
		213					Unspecified R
		214					Unspecified R
		215					Unspecified R
		216					Unspecified R
		217					Unspecified R
		218					Unspecified R
		219					Unspecified R
		220					Unspecified R
		221					Unspecified R
		222					Unspecified R
		223					Unspecified R
		224					Unspecified R
		225					Unspecified R
		226					Unspecified R
		227					Unspecified R
		228					Unspecified R
		229					Unspecified R
		230					Unspecified R
		231					Unspecified R
		232					Unspecified R
		233					Unspecified R
		234					Unspecified R
		235					Unspecified R
		236					Unspecified R
		237					Unspecified R
		238					Unspecified R
		239					Unspecified R
		240					Unspecified R
		241					Unspecified R
		242					Unspecified R
		243					Unspecified R
		244					Unspecified R
		245					Unspecified R
		246					Unspecified R
		247					Unspecified R
		248					Unspecified R
		249					Unspecified R
		250					Unspecified R
		251					Unspecified R
		252					Unspecified R
		253					Unspecified R
		254					Unspecified R
		255					Unspecified R
		256					Unspecified R
		257					Unspecified R
		258					Unspecified R
		259					Unspecified R
		260					Unspecified R
		261					Unspecified R
		262					Unspecified R
		263					Unspecified R
		264					

Site 368 Hole		Core 37		Cored Interval: 531.5-541.0 m	
AGE	ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	LITHOLOGIC DESCRIPTION
			0	VOID	CLAYSTONE AND PORCELLANITE WITH MINOR SILT
			1	CLAYSTONE, dark gray (5G 4/1) with interbedded 5 cm bands of olive back (5Y 7/1), hard, slight disturbance, bioturbated, PORCELLANITE, medium blue gray (5B 5/1) scattered throughout.	CLAYSTONE, dark gray-green (5G 4/1) and dark greenish gray (5G 4/1) with interbedded 5 cm bands of olive back (5Y 7/1), hard, slight disturbance, bioturbated throughout, sandy burrow fillings and occasional SILT layers.
			2	CLAYSTONE, dark gray (5G 4/1)	Color banding is very complex throughout core. Lower boundary of darker unit always very sharp and usually the upper boundary is gradational.
			3	CLAYSTONE, dark gray (5G 4/1)	Burrowing is beautifully displayed in this core.
			4	CLAYSTONE, dark gray (5G 4/1)	PORCELLANITE becoming almost CHERT, olive black (5Y 2/1) usually occurring in the claystone of the same color.
			5	CLAYSTONE, dark gray (5G 4/1)	SS at 2-58 (minor lithology)
			6	CLAYSTONE, dark gray (5G 4/1)	Quartz R
			7	CLAYSTONE, dark gray (5G 4/1)	Volcanic glass R
			8	CLAYSTONE, dark gray (5G 4/1)	Zeolites R
			9	CLAYSTONE, dark gray (5G 4/1)	Rads R
			10	CLAYSTONE, dark gray (5G 4/1)	SS at 5-30 (dominant lithology)
			11	CLAYSTONE, dark gray (5G 4/1)	Quartz R
			12	CLAYSTONE, dark gray (5G 4/1)	Volcanic glass R
			13	CLAYSTONE, dark gray (5G 4/1)	Pyrite R
			14	CLAYSTONE, dark gray (5G 4/1)	Dolomite R
			15	CLAYSTONE, dark gray (5G 4/1)	Feldspar R
			16	CLAYSTONE, dark gray (5G 4/1)	Clay R
			17	CLAYSTONE, dark gray (5G 4/1)	Pyrite R
			18	CLAYSTONE, dark gray (5G 4/1)	Carbonate Bomb: 5-9 to 10 cm = 1%

Site 368 Hole		Core 35		Cored Interval: 484.0-493.5 m	
AGE	ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	LITHOLOGIC DESCRIPTION
			0	VOID	CLAYSTONE AND PORCELLANITE
			1	CLAYSTONE, greenish gray (5G 6/1) alternating with dark greenish gray (5G 4/1) interbeds, each about 5 cm thick, hard, slight drilling disturbance, bioturbated, PORCELLANITE, medium blue gray (5B 5/1) scattered throughout.	CLAYSTONE, greenish gray (5G 6/1) alternating with dark greenish gray (5G 4/1) interbeds, each about 5 cm thick, hard, slight drilling disturbance, bioturbated, PORCELLANITE, medium blue gray (5B 5/1) scattered throughout.
			2	CLAYSTONE, dark gray (5G 4/1)	Color boundaries have sharp lower contact of dark colors and very gradational upwards to another sharp dark boundary.
			3	CLAYSTONE, dark gray (5G 4/1)	SILT, thin interbeds and burrow fillings occur scattered throughout core.
			4	CLAYSTONE, dark gray (5G 4/1)	PORCELLANITE, medium bluish gray (5B 5/1) occurs interbedded throughout core.
			5	CLAYSTONE, dark gray (5G 4/1)	SILT, thin interbeds and burrow fillings occur scattered throughout core.
			6	CLAYSTONE, dark gray (5G 4/1)	SS at 2-40 (minor lithology)
			7	CLAYSTONE, dark gray (5G 4/1)	Quartz D
			8	CLAYSTONE, dark gray (5G 4/1)	Heavy minerals R
			9	CLAYSTONE, dark gray (5G 4/1)	Fe/Mn R
			10	CLAYSTONE, dark gray (5G 4/1)	SS at 3-80 (dominant lithology)
			11	CLAYSTONE, dark gray (5G 4/1)	Clay D
			12	CLAYSTONE, dark gray (5G 4/1)	Pyrite R
			13	CLAYSTONE, dark gray (5G 4/1)	Forams R
			14	CLAYSTONE, dark gray (5G 4/1)	SS at 3-110 (minor lithology)
			15	CLAYSTONE, dark gray (5G 4/1)	Feldspar R
			16	CLAYSTONE, dark gray (5G 4/1)	Carbonate D
			17	CLAYSTONE, dark gray (5G 4/1)	Dolomite unspecified R
			18	CLAYSTONE, dark gray (5G 4/1)	Pyrite R
			19	CLAYSTONE, dark gray (5G 4/1)	SS at 3-140 (minor lithology)
			20	CLAYSTONE, dark gray (5G 4/1)	Quartz R
			21	CLAYSTONE, dark gray (5G 4/1)	Volcanic glass R
			22	CLAYSTONE, dark gray (5G 4/1)	Zeolites R
			23	CLAYSTONE, dark gray (5G 4/1)	Carbonate Bomb: 3-105 to 107 cm = 1%
			24	CLAYSTONE, dark gray (5G 4/1)	Core Catcher

Site 368		Hole	Core 38	Cored Interval : 560.0-569.5 m
AGE	ZONES	Fossil Character	Fossils	Lithology
UPPER CRETACEOUS 7-LOWER EOCENE?				
0				CLAYSTONE AND PORCELLANITE WITH MINOR SILT
				CLAYSTONE, dark greenish gray (56Y 4/1) with interbeds of dark gray green (56 4/1), hard, slight drilling disturbance, bioturbated, burrows filled with silt occasionally, some silt layers occur. Color banding on a fine scale is present but indistinct.
				PORCELLANITE grading to CHERI, greenish black (56 2/1) occur scattered throughout the core.
				Carbonate Bomb: 3-80 to 81 cm = 0%
1	0.5	VOID		
	1.0	△		
2	1.0	△		84
	1.5	△		
	2.0	△		
	2.5	△		
	3.0	△		
	3.5	△		
	4.0	△		
	4.5	△		
	5.0	△		
	5.5	△		
	6.0	△		
	6.5	△		
	7.0	△		
	7.5	△		
	8.0	△		
	8.5	△		
	9.0	△		
	9.5	△		
	10.0	△		
	10.5	△		
	11.0	△		
	11.5	△		
	12.0	△		
	12.5	△		
	13.0	△		
	13.5	△		
	14.0	△		
	14.5	△		
	15.0	△		
	15.5	△		
	16.0	△		
	16.5	△		
	17.0	△		
	17.5	△		
	18.0	△		
	18.5	△		
	19.0	△		
	19.5	△		
	20.0	△		
	20.5	△		
	21.0	△		
	21.5	△		
	22.0	△		
	22.5	△		
	23.0	△		
	23.5	△		
	24.0	△		
	24.5	△		
	25.0	△		
	25.5	△		
	26.0	△		
	26.5	△		
	27.0	△		
	27.5	△		
	28.0	△		
	28.5	△		
	29.0	△		
	29.5	△		
	30.0	△		
	30.5	△		
	31.0	△		
	31.5	△		
	32.0	△		
	32.5	△		
	33.0	△		
	33.5	△		
	34.0	△		
	34.5	△		
	35.0	△		
	35.5	△		
	36.0	△		
	36.5	△		
	37.0	△		
	37.5	△		
	38.0	△		
	38.5	△		
	39.0	△		
	39.5	△		
	40.0	△		
	40.5	△		
	41.0	△		
	41.5	△		
	42.0	△		
	42.5	△		
	43.0	△		
	43.5	△		
	44.0	△		
	44.5	△		
	45.0	△		
	45.5	△		
	46.0	△		
	46.5	△		
	47.0	△		
	47.5	△		
	48.0	△		
	48.5	△		
	49.0	△		
	49.5	△		
	50.0	△		
	50.5	△		
	51.0	△		
	51.5	△		
	52.0	△		
	52.5	△		
	53.0	△		
	53.5	△		
	54.0	△		
	54.5	△		
	55.0	△		
	55.5	△		
	56.0	△		
	56.5	△		
	57.0	△		
	57.5	△		
	58.0	△		
	58.5	△		
	59.0	△		
	59.5	△		
	60.0	△		
	60.5	△		
	61.0	△		
	61.5	△		
	62.0	△		
	62.5	△		
	63.0	△		
	63.5	△		
	64.0	△		
	64.5	△		
	65.0	△		
	65.5	△		
	66.0	△		
	66.5	△		
	67.0	△		
	67.5	△		
	68.0	△		
	68.5	△		
	69.0	△		
	69.5	△		
	70.0	△		
	70.5	△		
	71.0	△		
	71.5	△		
	72.0	△		
	72.5	△		
	73.0	△		
	73.5	△		
	74.0	△		
	74.5	△		
	75.0	△		
	75.5	△		
	76.0	△		
	76.5	△		
	77.0	△		
	77.5	△		
	78.0	△		
	78.5	△		
	79.0	△		
	79.5	△		
	80.0	△		
	80.5	△		
	81.0	△		
	81.5	△		
	82.0	△		
	82.5	△		
	83.0	△		
	83.5	△		
	84.0	△		
	84.5	△		
	85.0	△		
	85.5	△		
	86.0	△		
	86.5	△		
	87.0	△		
	87.5	△		
	88.0	△		
	88.5	△		
	89.0	△		
	89.5	△		
	90.0	△		
	90.5	△		
	91.0	△		
	91.5	△		
	92.0	△		
	92.5	△		
	93.0	△		
	93.5	△		
	94.0	△		
	94.5	△		
	95.0	△		
	95.5	△		
	96.0	△		
	96.5	△		
	97.0	△		
	97.5	△		
	98.0	△		
	98.5	△		
	99.0	△		
	99.5	△		
	100.0	△		
	100.5	△		
	101.0	△		
	101.5	△		
	102.0	△		
	102.5	△		
	103.0	△		
	103.5	△		
	104.0	△		
	104.5	△		
	105.0	△		
	105.5	△		
	106.0	△		
	106.5	△		
	107.0	△		
	107.5	△		
	108.0	△		
	108.5	△		
	109.0	△		
	109.5	△		
	110.0	△		
	110.5	△		
	111.0	△		
	111.5	△		
	112.0	△		
	112.5	△		
	113.0	△		
	113.5	△		
	114.0	△		
	114.5	△		
	115.0	△		
	115.5	△		
	116.0	△		
	116.5	△		
	117.0	△		
	117.5	△		
	118.0	△		
	118.5	△		
	119.0	△		
	119.5	△		
	120.0	△		
	120.5	△		
	121.0	△		
	121.5	△		
	122.0	△		
	122.5	△		
	123.0	△		
	123.5	△		
	124.0	△		
	124.5	△		
	125.0	△		
	125.5	△		
	126.0	△		
	126.5	△		
	127.0	△		
	127.5	△		
	128.0	△		
	128.5	△		
	129.0	△		
	129.5	△		
	130.0	△		
	130.5	△		
	131.0	△		
	131.5	△		
	132.0	△		
	132.5	△		
	133.0	△		
	133.5	△		
	134.0	△		
	134.5	△		
	135.0	△		
	135.5	△		
	136.0	△		
	136.5	△		
	137.0	△		
	137.5	△		
	138.0	△		
	138.5	△		
	139.0	△		
	139.5	△		
	140.0	△		
	140.5	△		
	141.0	△		
	141.5	△		
	142.0	△		
	142.5	△		
	143.0	△		
	143.5	△		
	144.0	△		
	144.5	△		
	145.0	△		
	145.5	△		
	146.0	△		
	146.5	△		
	147.0	△		
	147.5	△		
	148.0	△		
	148.5	△		
	149.0	△		
	149.5	△		
	150.0	△		
	150.5	△		
	151.0	△		
	151.5	△		
	152.0	△		
	152.5	△		
	153.0	△		
	153.5	△		
	154.0	△		
	154.5	△		
	155.0	△		
	155.5	△		
	156.0	△		
	156.5	△		
	157.0	△		
	157.5	△		
	158.0	△		
	158.5	△		
	159.0	△		
	159.5	△		
	160.0	△		
	160.5	△		
	161.0	△		
	161.5	△		
	162.0	△		
	162.5	△		
	163.0	△		
	163.5	△		
	164.0	△		
	164.5	△		
	165.0	△		
	165.5	△		
	166.0	△		
	166.5	△		
	167.0	△		
	167.5	△		
	168.0	△		
	168.5	△		
	169.0	△		
	169.5	△		
	170.			

Site 368		Core 40	Cored Interval : 579.0-588.5 m		
Age	ZONES	Fossil Character	Meter	Lithology	Lithologic Description
			0	VOID	CLAYSTONE SILT AND CHERIT CLAYSTONE, olive gray (SY 4/1) with olive black (SY 2/1) and greenish gray (SY 6/1) thin interbeds throughout, hard, slight drilling disturbance, bioturbated. Thin SILT layers common throughout, generally in lower portion of dark layers, upper and lower contacts sharp. A 14 cm thick bed of DOLOMITE or SIDERITE occurs at 4-32 to 146.
			0.5		CHERIT, olive black (SY 2/1) occurs scattered throughout as burrow filling and as discrete beds.
			1		Whole core is laminated.
			1.0		SS at 2-123 (minor lithology) Quartz A Carbonate Mica R unspecified Heavy minerals R F siderite Clay A Phosphates Pyrite R
			2		SS at 4-100 (dominant lithology) Quartz Clay Heavy minerals R Zeolites SS at 4-160 (minor lithology) Siderite D
			3		Carbonate Bomb : 4-141 to 142 cm = 81%
			4		
					100
					Core Catcher
					N R F R

3

Site 368 Hole		Core 43	Cored Interval : 607.5-617.0 m
			LITHOLOGIC DESCRIPTION
AGE	FORMAMS	RADS	LITHO. SAMPLE
ZONES	FOSSIL CHARACTER	PRES.	DEFORMATION
METERS	SECTION	SECT. 0	LITHOLOGY
0	CLAYSTONE AND PORCELLANITE WITH MINOR SILT	0.5	VOID
1	CLAYSTONE, olive black (5Y 2/1) with thin (2 to 5 cm) interbeds of olive gray (5Y 4/1), hard, no drilling disturbance, bioturbated, fine wavy laminations, rare thin (1 to 2 cm) SILT beds.	1.0	CLAYSTONE, olive black (5Y 2/1) interbedded with olive gray (5Y 4/1), hard, no drilling disturbance, bioturbated, sparse wavy laminations, a few SILT zones (especially in Section 4).
2	PORCELLANITE, occurs in darker unit.	2	PORCELLANITE, thin burrow fillings and laminae scattered throughout core.
3	SS at 2-88 (dominant lithology) Quartz R Pyrite R Heavy minerals R Clay R Zeolites D Radis R	3	SS at 3-91 (minor lithology) Clay D Radis R
4	SS at 2-92 (minor lithology) Quartz R Pyrite R Fish debris R	4	Carbonate Bomb : 4-68 to 69 cm = 3%
	Carbone Bomb : 3-93 to 94 cm = 10%		
			Core Catcher
		N R F	— — —

UPPER CRETACEOUS-LOWER EOCENE?

Site 368 Hole		Core 42	Cored Interval : 598.0-607.5 m
			LITHOLOGIC DESCRIPTION
AGE	FORMAMS	RADS	LITHO. SAMPLE
ZONES	FOSSIL CHARACTER	PRES.	DEFORMATION
METERS	SECTION	SECT. 0	LITHOLOGY
0	CLAYSTONE AND PORCELLANITE WITH MINOR SILT	0.5	VOID
1	CLAYSTONE, olive black (5Y 2/1) with thin (2 to 5 cm) interbeds of olive gray (5Y 4/1), hard, no drilling disturbance, bioturbated, fine wavy laminations, rare thin (1 to 2 cm) SILT beds.	1.0	CLAYSTONE AND PORCELLANITE WITH MINOR SILT
2	SS at 2-88 (dominant lithology) Quartz R Pyrite R Heavy minerals R Clay R Zeolites D Radis R	88	SS at 2-92 (minor lithology) Quartz R Pyrite R Fish debris R
3	SS at 2-92 (minor lithology) Quartz R Pyrite R Fish debris R	92	Carbone Bomb : 3-93 to 94 cm = 10%
4	SS at 2-92 (minor lithology) Quartz R Pyrite R Fish debris R	91	
			Core Catcher
		N R F	— — —

UPPER CRETACEOUS-LOWER EOCENE?

Site 368 Hole		Core 44	Cored Interval: 617.0-626.5 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY
UPPER CRETACEOUS? - LOWER EOCENE?			
Site 368 Hole		Core 45	Cored Interval: 626.5-636.0 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY
UPPER CRETACEOUS? - LOWER EOCENE?			

Site 368 Hole		Core 45	Cored Interval: 626.5-636.0 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY
UPPER CRETACEOUS? - LOWER EOCENE?			

D

Site 368 Hole Core 47 Cored Interval: 645.5-655.0 m								
LITHOLOGIC DESCRIPTION								
LITHO. SAMPLE	DEFORMATION	LITHOLOGY	METERS	SECTION	FOSIL	ZONES	FOSSIL CHARACTER	AGE
		INTERBEDDED CLAYSTONE AND SILTSTONE CLAYSTONE, olive black (SY 2/1) interbedded with dark greenish gray (SY 4/1), hard, no drilling disturbance, bioturbated, basal contact of drier bed sharp, upper contact gradational, fine wavy laminations. Pyrite rosettes occur in Sections 3 and 4.	0					
		SILTSTONE, olive black (SY 2/1), hard, occur as 1 to 5 cm thick beds scattered throughout core - sometimes making up over 50% of a section (i.e. Section 3). Thickness of interbeds vary between 1 and 40 cm.	0.5					
		Thickness of interbeds vary between 1 and 40 cm.	1					
		Carbonate Bomb: 3-146 to 147 cm = 0%	1.0					
			2					
			3					
			4					
			5					
			6					
			85					
UPPER CRETACEOUS-LOWER EOCENE?								
Site 368 Hole Core 46 Cored Interval: 636.0-645.5 m								
LITHOLOGIC DESCRIPTION								
LITHO. SAMPLE	DEFORMATION	LITHOLOGY	METERS	SECTION	FOSIL	ZONES	FOSSIL CHARACTER	AGE
		INTERBEDDED CLAYSTONE AND SILTSTONE CLAYSTONE, olive black (SY 2/1) and olive gray (SY 4/1), hard, no drilling disturbance, bioturbated, basal contact of drier bed sharp, upper contact gradational, fine wavy laminations. Pyrite rosettes occur in Sections 3 and 4.	0					
		SILTSTONE, olive black (SY 2/1), hard, occur as 1 to 5 cm thick beds scattered throughout core - sometimes making up over 50% of a section (i.e. Section 3).	0.5					
		Thickness of interbeds vary between 1 and 40 cm.	1					
		Carbonate Bomb: 3-146 to 147 cm = 0%	1.0					
			2					
			3					
			4					
			5					
			6					
			85					
UPPER CRETACEOUS-LOWER EOCENE?								

Site 368, Core 48, Cored Interval: 664.5-674.0 m: NO RECOVERY
Site 368, Core 49, Cored Interval: 693.0-702.5 m: NO RECOVERY

103

Site 368		Core 61		Cored Interval: 961.0-968.5 m	
ZONES	FORMAMS	FORMAMS	NAMESS	FOSSTIL	FOSSTIL
AGE				CHARACTER	CHARACTER
				PPLD.	PPLS.
				SECTION	METERS
				LITHOLOGY	DEFORMATION
				LITHO. SAMPLE	LITHOLOGIC DESCRIPTION
				Olivine Diabase, medium gray (N5), coarse euhedral texture.	

Site 368 Hole		Core 62 Cored Interval: 968.5-975.0 m			
AGE	ZONES	CHARACTER	FOSIL	SECTORS	LITHOLOGY
FORMAMS	NAME	ABUND.	PRES.	METERS	LITHO. SAMPLE
				0	OILY DIABASE AND SHALE
				0.5	OLIVE DIABASE, medium gray (NS), coarse euhedral texture becoming finer with depth.
				1.0	Baked contact zone of SHALE, medium light gray (N6), light brownish gray (5R 6/1) to dark gray (N3), very hard and fissile, baked zone extends from 3-7 cm to 100 cm, burrowed in unbaked portion.
				1.5	SS at 4-36 (dominant lithology)
				2.0	Quartz C Pyrite C Nannos R Volcanic glass R
				2.5	
				3.0	
				3.5	
				4.0	
				4.5	
				5.0	
				5.5	
				6.0	
				6.5	
				7.0	
				7.5	
				8.0	
				8.5	
				9.0	
				9.5	
				10.0	
				10.5	
				11.0	
				11.5	
				12.0	
				12.5	
				13.0	
				13.5	
				14.0	
				14.5	
				15.0	
				15.5	
				16.0	
				16.5	
				17.0	
				17.5	
				18.0	
				18.5	
				19.0	
				19.5	
				20.0	
				20.5	
				21.0	
				21.5	
				22.0	
				22.5	
				23.0	
				23.5	
				24.0	
				24.5	
				25.0	
				25.5	
				26.0	
				26.5	
				27.0	
				27.5	
				28.0	
				28.5	
				29.0	
				29.5	
				30.0	
				30.5	
				31.0	
				31.5	
				32.0	
				32.5	
				33.0	
				33.5	
				34.0	
				34.5	
				35.0	
				35.5	
				36.0	
				36.5	
				37.0	
				37.5	
				38.0	
				38.5	
				39.0	
				39.5	
				40.0	
				40.5	
				41.0	
				41.5	
				42.0	
				42.5	
				43.0	
				43.5	
				44.0	
				44.5	
				45.0	
				45.5	
				46.0	
				46.5	
				47.0	
				47.5	
				48.0	
				48.5	
				49.0	
				49.5	
				50.0	
				50.5	
				51.0	
				51.5	
				52.0	
				52.5	
				53.0	
				53.5	
				54.0	
				54.5	
				55.0	
				55.5	
				56.0	
				56.5	
				57.0	
				57.5	
				58.0	
				58.5	
				59.0	
				59.5	
				60.0	
				60.5	
				61.0	
				61.5	
				62.0	
				62.5	
				63.0	
				63.5	
				64.0	
				64.5	
				65.0	
				65.5	
				66.0	
				66.5	
				67.0	
				67.5	
				68.0	
				68.5	
				69.0	
				69.5	
				70.0	
				70.5	
				71.0	
				71.5	
				72.0	
				72.5	
				73.0	
				73.5	
				74.0	
				74.5	
				75.0	
				75.5	
				76.0	
				76.5	
				77.0	
				77.5	
				78.0	
				78.5	
				79.0	
				79.5	
				80.0	
				80.5	
				81.0	
				81.5	
				82.0	
				82.5	
				83.0	
				83.5	
				84.0	
				84.5	
				85.0	
				85.5	
				86.0	
				86.5	
				87.0	
				87.5	
				88.0	
				88.5	
				89.0	
				89.5	
				90.0	
				90.5	
				91.0	
				91.5	
				92.0	
				92.5	
				93.0	
				93.5	
				94.0	
				94.5	
				95.0	
				95.5	
				96.0	
				96.5	
				97.0	
				97.5	
				98.0	
				98.5	
				99.0	
				99.5	
				100.0	
				100.5	
				101.0	
				101.5	
				102.0	
				102.5	
				103.0	
				103.5	
				104.0	
				104.5	
				105.0	
				105.5	
				106.0	
				106.5	
				107.0	
				107.5	
				108.0	
				108.5	
				109.0	
				109.5	
				110.0	
				110.5	
				111.0	
				111.5	
				112.0	
				112.5	
				113.0	
				113.5	
				114.0	
				114.5	
				115.0	
				115.5	
				116.0	
				116.5	
				117.0	
				117.5	
				118.0	
				118.5	
				119.0	
				119.5	
				120.0	
				120.5	
				121.0	
				121.5	
				122.0	
				122.5	
				123.0	
				123.5	
				124.0	
				124.5	
				125.0	
				125.5	
				126.0	
				126.5	
				127.0	
				127.5	
				128.0	
				128.5	
				129.0	
				129.5	
				130.0	
				130.5	
				131.0	
				131.5	
				132.0	
				132.5	
				133.0	
				133.5	
				134.0	
				134.5	
				135.0	
				135.5	
				136.0	
				136.5	
				137.0	
				137.5	
				138.0	
				138.5	
				139.0	
				139.5	
				140.0	
				140.5	
				141.0	
				141.5	
				142.0	
				142.5	
				143.0	
				143.5	
				144.0	
				144.5	
				145.0	
				145.5	
				146.0	
				146.5	
				147.0	
				147.5	
				148.0	
				148.5	
				149.0	
				149.5	
				150.0	
				150.5	
				151.0	
				151.5	
				152.0	
				152.5	
				153.0	
				153.5	
				154.0	
				154.5	
				155.0	
				155.5	
				156.0	
				156.5	
				157.0	
				157.5	
				158.0	
				158.5	
				159.0	
				159.5	
				160.0	
				160.5	
				161.0	
				161.5	
				162.0	
				162.5	
				163.0	
				163.5	
				164.0	
				164.5	
				165.0	
				165.5	
				166.0	
				166.5	
				167.0	
				167.5	
				168.0	
				168.5	
				169.0	
				169.5	
				170.0	
				170.5	
				171.0	
				171.5	
				172.0	
				172.5	
				173.0	
				173.5	
				174.0	
				174.5	
				1	

109
DEEP SEA DRILLING PROJECT

LEG 41 SITE 369 (HOLE 369A)

SITE SUMMARY SHEET

PRINCIPAL RESULTS:

Continental Slope off Spanish Sahara

The upper part of the sediment section of the continental slope off Cape Bojador was continuously cored down to 488.5 meters. The sediments are predominantly pelagic with only a minor coarse terrigenous contribution. They consist mainly of nannofossil marls, with some limestone and chalk. The section can be summarized as follows (from top to bottom); 1) Quaternary to early Miocene nannofossil marl and ooze with minor volcanic ash; 2) early Miocene to middle Eocene siliceous nannofossil marl; 3) middle Eocene to late Cretaceous argillaceous nannofossil limestone and chalk with occasional chert; and 4) late Cretaceous to late Aptian dark-colored silty nannofossil marls.

No shallow water material was observed and redeposition by slumping appears to be relatively minor in importance. The stratigraphy seems to be relatively continuous but minor hiatuses were found in the late Pliocene and early late Eocene, and larger ones affect the early Eocene, Paleocene and Turonian to late Albian. Water depth conditions do not show appreciable variations and the sediments reflect a continental slope environment above the CCD since the late Aptian. The dark brown to blackish Aptian-Albian silty marls contain barite rosettes and layers partially replaced by calcite, possibly resulting from the upward and/or lateral migration of solutions. A strong lower reflector is of middle to late Eocene age and corresponds with an angular unconformity. Another reflector marks the passage from relatively carbonate-poor to carbonate-rich sediments during the early Miocene. The hole had to be abandoned for safety reasons when relatively low methane/ethane ratios in gas were encountered.

Date occupied	2335 March 23, 1975
Date departed	2050 March 26, 1975
Time on site	2 days, 21 hrs, 15 minutes
Position: Latitude	26°35.5'N
Longitude	14°59.9'W
Water depth (sea level)	1752 corrected meters
Water depth (rig floor)	1762 corrected meters

LEG 41 SITE 369 (HOLE 369A)
SITE SUMMARY SHEET, con't.

Penetration	488.5 meters
Number of holes	2
Number of cores	52
Total length of cored section	488.5 meters
Total core recovered	386.4 meters
Percentage core recovery	79%


Oldest Sediment Cored

Depth subbottom	488.5 meters
Nature	Silty marl
Age	Aptian
Measured velocity	1.9 km/sec

Site 369 Hole		Core 1	Cored Interval : 0.0-4.0 m
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGY
METERS	SECTORS	PRES.	DEFORMATION
0	F A G R R M N A G	void	10YR 7/1
0.5			134
1			144
2			56Y 3/1
100			
			SS at 2-100 (dominant lithology)
			Quartz R Clay C
			Glaucite R Fe/Mn R
			Carbonate rhombs R Carbonate rhombs R
			unspecified C Forams C
			Nannos A Radls R
			Sponge spicules R Silicoflagellates R
			Fish debris R Pollen R
			SS at 1-144 (minor lithology)
			Quartz R Clay C
			Glaucite R Fe/Mn R
			Carbonate rhombs R Carbonate rhombs R
			unspecified C Forams C
			Nannos A Radls R
			Silicoflagellates R
			Fish debris R Pollen R
			SS at 1-133 (minor lithology)
			Quartz R Clay C
			Glaucite R Fe/Mn R
			Carbonate rhombs R Carbonate rhombs R
			Forams C
			Nannos A Radls R
			Diatoms R Fish debris R
			SS at 1-133 (minor lithology)
			Quartz R Clay C
			Glaucite R Fe/Mn R
			Carbonate rhombs R Carbonate rhombs R
			Forams C
			Nannos A Radls R
			Diatoms R Fish debris R

Site 369 Hole		Core 2	Cored Interval : 4.0-13.5 m
AGE	ZONES	FOSSIL CHARACTER	LITHOLOGIC DESCRIPTION
METERS	SECTORS	PRES.	LITHO. SAMPLE
0	F A G R R P		CLAYEY NANNO Ooze, greenish gray (5GY 6/1), with top 50 cm grayish orange (10YR 7/4) and greenish gray (5GY 6/1), soft, severe drilling disturbance, black streaks and blebs throughout, rare bioturbation apparent.
1			
1.0			
1.5			
2			
3			
4			
5			
6			
			DISCOASSTER ASYMMETRICUS-RETICULOFENESTRA PSEUDOMORBILLICA (NN14-NN15)
			DISCOASSTER GLABROROTATILA MARGARITATE EVOLOTA (NN14-NN15)
			GLABROROTATILA MARGARITATE MARGINIFERAE (NN14-NN15)

112



114

Site 369		Hole A	Cone 7	Cored interval: 99-0-108.5 m
ZONES	AGE	FOSIL	LITHOLOGY	LITHOLOGIC DESCRIPTION
MIDDLE MIOCENE				
Sphenolithus heteromorphus NNS				
5				
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
24				
25				
26				
27				
28				
29				
30				
31				
32				
33				
34				
35				
36				
37				
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				
48				
49				
50				
51				
52				
53				
54				
55				
56				
57				
58				
59				
60				
61				
62				
63				
64				
65				
66				
67				
68				
69				
70				
71				
72				
73				
74				
75				
76				
77				
78				
79				
80				
81				
82				
83				
84				
85				
86				
87				
88				
89				
90				
91				
92				
93				
94				
95				
96				
97				
98				
99				
100				
101				
102				
103				
104				
105				
106				
107				
108				
109				
110				
111				
112				
113				
114				
115				
116				
117				
118				
119				
120				
121				
122				
123				
124				
125				
126				
127				
128				
129				
130				
131				
132				
133				
134				
135				
136				
137				
138				
139				
140				
141				
142				
143				
144				
145				
146				
147				
148				
149				
150				
151				
152				
153				
154				
155				
156				
157				
158				
159				
160				
161				
162				
163				
164				
165				
166				
167				
168				
169				
170				
171				
172				
173				
174				
175				
176				
177				
178				
179				
180				
181				
182				
183				
184				
185				
186				
187				
188				
189				
190				
191				
192				
193				
194				
195				
196				
197				
198				
199				
200				
201				
202				
203				
204				
205				
206				
207				
208				
209				
210				
211				
212				
213				
214				
215				
216				
217				
218				
219				
220				
221				
222				
223				
224				
225				
226				
227				
228				
229				
230				
231				
232				
233				
234				
235				
236				
237				
238				
239				
240				
241				
242				
243				
244				
245				
246				
247				
248				
249				
250				
251				
252				
253				
254				
255				
256				
257				
258				
259				
260				
261				
262				
263				
264				
265				
266				
267				
268				
269				
270				
271				
272				
273				
274				
275				
276				
277				
278				
279				
280				
281				
282				
283				
284				
285				
286				
287				
288				
289				
290				
291				
292				
293				
294				
295				
296				
297				
298				
299				
300				
301				
302				
303				
304				
305				
306				
307				
308				
309				
310				
311				
312				
313				
314				
315				
316				
317				
318				
319				
320				
321				
322				
323				
324				
325				
326				
327				
328				
329				
330				
331				
332				
333				
334				
335				
336				
337				
338				
339				
340				
341				
342				
343				
344				
345				
346				
347				
348				
349				
350				
351				
352				
353				
354				
355				
356				
357				
358				
359				
360				
361				
362				
363				
364				
365				
366				
367				
368				
369				
370				
371				
372				
373				
374				
375				
376				
377				
378				
379				
380				
381				
382				
383				
384				
385				
386				
387				
388				
389				
390				
391				
392				
393				
394				
395				
396				
397				
398				
399				
400				
401				
402				
403				
404				
405				
406				
407				
408				
409				
410				
411				
412				
413				
414				
415				
416				
417				
418				
419				
420				
421				
422				
423				
424				
425				
426				
427				
428				
429				
430				
431				
432				
433				

Site 369 Hole A		Core 10	Cored Interval : 127.5-137.0 m										
AGE	ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	SECTION	SECTORS	ABUND.	FOSIL	ABUND.	PRAMS	DEFORMATION	LITHO. SAMPLE	LITHOLOGIC DESCRIPTION
			0	VOID	0.5								SS at 1-100 (dominant lithology)
			1		1.0	R	100						Quartz R Clay A Micronodules R Carbonate C Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			2										SS at 2-50 (minor lithology)
			3										Quartz C Clay R Heavy minerals A Fe/Mn R Micronodules A Carbonate C Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			4										SS at 3-80 (dominant lithology)
			5										Quartz R Clay R Heavy minerals R Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			6										SS at 4-50 (dominant lithology)
			7										Quartz R Clay R Heavy minerals R Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			8										SS at 5-60 (dominant lithology)
			9										Quartz R Clay R Heavy minerals R Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			10										Carbone Bomb : 3-56 to 57 cm = 31%
													Carbone Bomb : 3-95 to 96 cm = 30%

LOWER Miocene

Stichocystites dissimilis-Globigerinoides trilobatus
Discocaster trilobatus NNN
Globigerinoides dissimilis-Globigerinoides trilobatus
Heterostrophopora ampliapertura NNA
Praeobulina glomerosa
Dorcadopsisyrta alata

Site 369 Hole A		Core 11	Cored Interval : 137.0-146.5 m										
AGE	ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	SECTION	SECTORS	ABUND.	FOSIL	ABUND.	PRAMS	DEFORMATION	LITHO. SAMPLE	LITHOLOGIC DESCRIPTION
			0	N	0.5								SS at 1-6 (minor lithology)
			1	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R Sponge spicules R Silicoflagellates R
			2	N	1.0	D							SS at 1-130 (dominant lithology)
			3	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			4	N	1.0	D							SS at 1-30 (dominant lithology)
			5	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			6	N	1.0	D							SS at 3-30 (dominant lithology)
			7	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			8	N	1.0	D							SS at 3-50 (dominant lithology)
			9	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			10	N	1.0	D							SS at 3-50 (dominant lithology)
			11	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			12	N	1.0	D							SS at 3-50 (dominant lithology)
			13	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			14	N	1.0	D							SS at 3-50 (dominant lithology)
			15	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			16	N	1.0	D							SS at 3-50 (dominant lithology)
			17	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			18	N	1.0	D							SS at 3-50 (dominant lithology)
			19	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			20	N	1.0	D							SS at 3-50 (dominant lithology)
			21	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			22	N	1.0	D							SS at 3-50 (dominant lithology)
			23	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			24	N	1.0	D							SS at 3-50 (dominant lithology)
			25	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			26	N	1.0	D							SS at 3-50 (dominant lithology)
			27	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			28	N	1.0	D							SS at 3-50 (dominant lithology)
			29	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			30	N	1.0	D							SS at 3-50 (dominant lithology)
			31	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			32	N	1.0	D							SS at 3-50 (dominant lithology)
			33	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			34	N	1.0	D							SS at 3-50 (dominant lithology)
			35	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			36	N	1.0	D							SS at 3-50 (dominant lithology)
			37	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			38	N	1.0	D							SS at 3-50 (dominant lithology)
			39	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			40	N	1.0	D							SS at 3-50 (dominant lithology)
			41	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			42	N	1.0	D							SS at 3-50 (dominant lithology)
			43	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			44	N	1.0	D							SS at 3-50 (dominant lithology)
			45	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			46	N	1.0	D							SS at 3-50 (dominant lithology)
			47	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			48	N	1.0	D							SS at 3-50 (dominant lithology)
			49	N	1.0	D							Quartz R Clay A Micronodules R Carbonate R Forams R Nannos R Diatoms A Rads R Fish debris R
			50	N	1.0	D							SS at 3-50 (dominant lithology)

Core 12		Cored Interval: 146.5-156.0 m			
Hole A	Site 369	Fossil Character	Lithology	Lithologic Description	
ZONES	AGE	Fossils	METERS	DEFORMATION	LITHO. SAMPLE
FORAMS		RADS.	0	Lx	NANO DIATOM MARL, olive gray (SY 3/2), firm, drilling breccia to moderate drilling disturbance, bioturbated, becomes AD-BEARING by Sect on 2.
NAOMIS		RADS.	0.5	D	SS at 2-70 (dominant lithology)
ABUND.		FOSILS	1	L	Quartz R
		PPRS.	1.0	N	Clay A
			2	D	Fe/Mn R
			3	D	Nanom A
			4	D	Diatoms A
			5	D	Silicoflage llates R
			6	D	Rads A
			7	D	Pollen R
			70		Fish debris R
					Pollen R
					Carboante Bomb: 3-77 to 78 cm = 14%
Diatoceraspis terebrapera Globigerinata dissimilis-Globigerinata staminiferae Dissimilis testudinaria dissimilis					
N F-C-P-H R C G Core F C M/G NS/G F Catcher					

Site 369 Hole A		Core 14 Cored Interval: 166.5-175.0 m	
		LITHOLOGIC DESCRIPTION	
ZONES	FOSIL CHARACTER	METERS	LITHOLOGY
ACE		0	NANNO DIATOM MARL, olive gray (SY 3/2), very firm, slight drilling disturbance except for drilling breccia at top 80 cm, bioturbated, yellowish flecks scattered throughout (may be mollusk fragments). A thin silt/sand layer may be a slump.
FORAMS	RADS	0.5	SS at 1.5-20 (dominant lithology)
MAMS		1	Quartz
		1.0	Clay
		1.5	Micronodules
		2	Mollusk debris
		3	Forams
		4	N
		5	N
		6	N
		50	
UPPER OLIGOCENE			
Sphenolithus cf perakensis NP25 Dolcadospyris astucinus?			
N C-F P R + P Core Catcher			

Site 369 Hole A		Core 15 Cored Interval: 175.0-184.5 m	
		LITHOLOGIC DESCRIPTION	
ZONES	FOSIL CHARACTER	METERS	LITHOLOGY
ACE		0	NANNO MARL, olive gray (SY 3/2), slight drilling disturbance, very firm, bioturbated, light yellow flecks scattered throughout (may be mollusk fragments). A fossil nassa zone occurs at 5-52 to 66.
FORAMS	RADS	0.5	SS at 2-100 (dominant lithology)
MAMS		1	Quartz
		1.0	Clay
		1.5	Micronodules
		2	Zeolites
		3	Forams
		4	Dolatons
		5	Rads
		6	N
		100	
UPPER OLIGOCENE			
Sphenolithus cf perakensis NP25 Dolcadospyris astucinus?			
N C-F P R + P Core Catcher			

Site 369 Hole A		Core 17		Cored Interval : 194.0-203.5 m	
ZONES	FOSSILS	CHARACTERS	LITHOLOGY	DEFORMATION	LITHO. SAMPLE
NP24	AG6	1 0.5 1.0 0	N N 0	10Y 4/2	
NP25	AB5UW	2 0	N 0	10Y 6/2	
NP26	FOS511	3 0	N N	55Y 5/2 100	
NP27	RADS	4 0	N 0	10Y 6/2 10Y 4/2	
NP28	METERS	5 0	N N	10Y 6/2	
NP29	SECTIONS	6 0	N D	5G 4/1	
NP30	PPS5			10Y 6/2	
NP31	ABSUD			10Y 6/2	
NP32	FOS511			10Y 6/2	
NP33	RADS			10Y 6/2	
NP34	METERS			10Y 6/2	
NP35	SECTIONS			10Y 6/2	
NP36	PPS5			10Y 6/2	
NP37	ABSUD			10Y 6/2	
NP38	FOS511			10Y 6/2	
NP39	RADS			10Y 6/2	
NP40	METERS			10Y 6/2	
NP41	SECTIONS			10Y 6/2	
NP42	PPS5			10Y 6/2	
NP43	ABSUD			10Y 6/2	
NP44	FOS511			10Y 6/2	
NP45	RADS			10Y 6/2	
NP46	METERS			10Y 6/2	
NP47	SECTIONS			10Y 6/2	
NP48	PPS5			10Y 6/2	
NP49	ABSUD			10Y 6/2	
NP50	FOS511			10Y 6/2	
NP51	RADS			10Y 6/2	
NP52	METERS			10Y 6/2	
NP53	SECTIONS			10Y 6/2	
NP54	PPS5			10Y 6/2	
NP55	ABSUD			10Y 6/2	
NP56	FOS511			10Y 6/2	
NP57	RADS			10Y 6/2	
NP58	METERS			10Y 6/2	
NP59	SECTIONS			10Y 6/2	
NP60	PPS5			10Y 6/2	
NP61	ABSUD			10Y 6/2	
NP62	FOS511			10Y 6/2	
NP63	RADS			10Y 6/2	
NP64	METERS			10Y 6/2	
NP65	SECTIONS			10Y 6/2	
NP66	PPS5			10Y 6/2	
NP67	ABSUD			10Y 6/2	
NP68	FOS511			10Y 6/2	
NP69	RADS			10Y 6/2	
NP70	METERS			10Y 6/2	
NP71	SECTIONS			10Y 6/2	
NP72	PPS5			10Y 6/2	
NP73	ABSUD			10Y 6/2	
NP74	FOS511			10Y 6/2	
NP75	RADS			10Y 6/2	
NP76	METERS			10Y 6/2	
NP77	SECTIONS			10Y 6/2	
NP78	PPS5			10Y 6/2	
NP79	ABSUD			10Y 6/2	
NP80	FOS511			10Y 6/2	
NP81	RADS			10Y 6/2	
NP82	METERS			10Y 6/2	
NP83	SECTIONS			10Y 6/2	
NP84	PPS5			10Y 6/2	
NP85	ABSUD			10Y 6/2	
NP86	FOS511			10Y 6/2	
NP87	RADS			10Y 6/2	
NP88	METERS			10Y 6/2	
NP89	SECTIONS			10Y 6/2	
NP90	PPS5			10Y 6/2	
NP91	ABSUD			10Y 6/2	
NP92	FOS511			10Y 6/2	
NP93	RADS			10Y 6/2	
NP94	METERS			10Y 6/2	
NP95	SECTIONS			10Y 6/2	
NP96	PPS5			10Y 6/2	
NP97	ABSUD			10Y 6/2	
NP98	FOS511			10Y 6/2	
NP99	RADS			10Y 6/2	
NP100	METERS			10Y 6/2	
NP101	SECTIONS			10Y 6/2	
NP102	PPS5			10Y 6/2	
NP103	ABSUD			10Y 6/2	
NP104	FOS511			10Y 6/2	
NP105	RADS			10Y 6/2	
NP106	METERS			10Y 6/2	
NP107	SECTIONS			10Y 6/2	
NP108	PPS5			10Y 6/2	
NP109	ABSUD			10Y 6/2	
NP110	FOS511			10Y 6/2	
NP111	RADS			10Y 6/2	
NP112	METERS			10Y 6/2	
NP113	SECTIONS			10Y 6/2	
NP114	PPS5			10Y 6/2	
NP115	ABSUD			10Y 6/2	
NP116	FOS511			10Y 6/2	
NP117	RADS			10Y 6/2	
NP118	METERS			10Y 6/2	
NP119	SECTIONS			10Y 6/2	
NP120	PPS5			10Y 6/2	
NP121	ABSUD			10Y 6/2	
NP122	FOS511			10Y 6/2	
NP123	RADS			10Y 6/2	
NP124	METERS			10Y 6/2	
NP125	SECTIONS			10Y 6/2	
NP126	PPS5			10Y 6/2	
NP127	ABSUD			10Y 6/2	
NP128	FOS511			10Y 6/2	
NP129	RADS			10Y 6/2	
NP130	METERS			10Y 6/2	
NP131	SECTIONS			10Y 6/2	
NP132	PPS5			10Y 6/2	
NP133	ABSUD			10Y 6/2	
NP134	FOS511			10Y 6/2	
NP135	RADS			10Y 6/2	
NP136	METERS			10Y 6/2	
NP137	SECTIONS			10Y 6/2	
NP138	PPS5			10Y 6/2	
NP139	ABSUD			10Y 6/2	
NP140	FOS511			10Y 6/2	
NP141	RADS			10Y 6/2	
NP142	METERS			10Y 6/2	
NP143	SECTIONS			10Y 6/2	
NP144	PPS5			10Y 6/2	
NP145	ABSUD			10Y 6/2	
NP146	FOS511			10Y 6/2	
NP147	RADS			10Y 6/2	
NP148	METERS			10Y 6/2	
NP149	SECTIONS			10Y 6/2	
NP150	PPS5			10Y 6/2	
NP151	ABSUD			10Y 6/2	
NP152	FOS511			10Y 6/2	
NP153	RADS			10Y 6/2	
NP154	METERS			10Y 6/2	
NP155	SECTIONS			10Y 6/2	
NP156	PPS5			10Y 6/2	
NP157	ABSUD			10Y 6/2	
NP158	FOS511			10Y 6/2	
NP159	RADS			10Y 6/2	
NP160	METERS			10Y 6/2	
NP161	SECTIONS			10Y 6/2	
NP162	PPS5			10Y 6/2	
NP163	ABSUD			10Y 6/2	
NP164	FOS511			10Y 6/2	
NP165	RADS			10Y 6/2	
NP166	METERS			10Y 6/2	
NP167	SECTIONS			10Y 6/2	
NP168	PPS5			10Y 6/2	
NP169	ABSUD			10Y 6/2	
NP170	FOS511			10Y 6/2	
NP171	RADS			10Y 6/2	
NP172	METERS			10Y 6/2	
NP173	SECTIONS			10Y 6/2	
NP174	PPS5			10Y 6/2	
NP175	ABSUD			10Y 6/2	
NP176	FOS511			10Y 6/2	
NP177	RADS			10Y 6/2	
NP178	METERS			10Y 6/2	
NP179	SECTIONS			10Y 6/2	
NP180	PPS5			10Y 6/2	
NP181	ABSUD			10Y 6/2	
NP182	FOS511			10Y 6/2	
NP183	RADS			10Y 6/2	
NP184	METERS			10Y 6/2	
NP185	SECTIONS			10Y 6/2	
NP186	PPS5			10Y 6/2	
NP187	ABSUD			10Y 6/2	
NP188	FOS511			10Y 6/2	
NP189	RADS			10Y 6/2	
NP190	METERS			10Y 6/2	
NP191	SECTIONS			10Y 6/2	
NP192	PPS5			10Y 6/2	
NP193	ABSUD			10Y 6/2	
NP194	FOS511			10Y 6/2	
NP195	RADS			10Y 6/2	
NP196	METERS			10Y 6/2	
NP197	SECTIONS			10Y 6/2	
NP198	PPS5			10Y 6/2	
NP199	ABSUD			10Y 6/2	
NP200	FOS511			10Y 6/2	
NP201	RADS			10Y 6/2	
NP202	METERS			10Y 6/2	
NP203	SECTIONS			10Y 6/2	
NP204	PPS5			10Y 6/2	
NP205	ABSUD			10Y 6/2	
NP206	FOS511			10Y 6/2	
NP207	RADS			10Y 6/2	
NP208	METERS			10Y 6/2	
NP209	SECTIONS			10Y 6/2	
NP210	PPS5			10Y 6/2	
NP211	ABSUD			10Y 6/2	
NP212	FOS511			10Y 6/2	
NP213	RADS			10Y 6/2	
NP214	METERS			10Y 6/2	
NP215	SECTIONS			10Y 6/2	
NP216	PPS5			10Y 6/2	
NP217	ABSUD			10Y 6/2	
NP218	FOS511			10Y 6/2	
NP219	RADS			10Y 6/2	
NP220	METERS			10Y 6/2	
NP221	SECTIONS			10Y 6/2	
NP222	PPS5			10Y 6/2	
NP223	ABSUD			10Y 6/2	
NP224	FOS511			10Y 6/2	
NP225	RADS			10Y 6/2	
NP226	METERS			10Y 6/2	
NP227	SECTIONS			10Y 6/2	
NP228	PPS5			10Y 6/2	
NP229	ABSUD			10Y 6/2	
NP230	FOS511			10Y 6/2	
NP231	RADS			10Y 6/2	
NP232	METERS			10Y 6/2	
NP233	SECTIONS			10Y 6/2	
NP234	PPS5			10Y 6/2	
NP235	ABSUD			10Y 6/2	
NP236	FOS511			10Y 6/2	
NP237	RADS			10Y 6/2	
NP238	METERS			10Y 6/2	
NP239	SECTIONS			10Y 6/2	
NP240	PPS5			10Y 6/2	
NP241	ABSUD			10Y 6/2	
NP242	FOS511			10Y 6/2	
NP243	RADS			10Y 6/2	
NP244	METERS			10Y 6/2	
NP245	SECTIONS			10Y 6/2	
NP246	PPS5			10Y 6/2	
NP247	ABSUD			10Y 6/2	
NP248	FOS511			10Y 6/2	
NP249	RADS			10Y 6/2	
NP250	METERS			10Y 6/2	
NP251	SECTIONS			10Y 6/2	
NP252	PPS5			10Y 6/2	
NP253	ABSUD			10Y 6/2	
NP254	FOS511			10Y 6/2	
NP255	RADS			10Y 6/2	
NP256	METERS			10Y 6/2	
NP257	SECTIONS			10Y 6/2	
NP258	PPS5			10Y 6/2	
NP259	ABSUD			10Y 6/2	
NP260	FOS511			10Y 6/2	
NP261	RADS			10Y 6/2	
NP262	METERS			10Y 6/2	
NP263	SECTIONS			10Y 6/2	
NP264	PPS5			10Y 6/2	
NP265	ABSUD			10Y 6/2	
NP266	FOS511			10Y 6/2	
NP267	RADS			10Y 6/2	
NP268	METERS			10Y 6/2	
NP269	SECTIONS			10Y 6/2	
NP270	PPS5			10Y 6/2	
NP271	ABSUD			10Y 6/2	
NP272	FOS511			10Y 6/2	
NP273	RADS			10Y 6/2	
NP274	METERS			10Y 6/2	
NP275	SECTIONS			10Y 6/2	
NP276	PPS5			10Y 6/2	
NP277	ABSUD			10Y 6/2	
NP278	FOS511			10Y 6/2	
NP279	RADS			10Y 6/2	
NP280	METERS			10Y 6/2	
NP281	SECTIONS			10Y 6/2	
NP282	PPS5			10Y 6/2	
NP283	ABSUD			10Y 6/2	
NP284	FOS511			10Y 6/2	
NP285	RADS			10Y 6/2	
NP286	METERS			10Y 6/2	
NP287	SECTIONS			10Y 6/2	
NP288	PPS5			10Y 6/2	
NP289	ABSUD			10Y 6/2	
NP290	FOS511			10Y 6/2	
NP291	RADS			10Y 6/2	
NP292	METERS</td				

Site 369		Hole A		Core 20		Cored Interval : 222.5-232.0 m	
ZONES	FOSIL CHARACTER	METERS	LITHOLOGY	SECTION	DEFORMATION	SAMPLE	LITHO. SAMPLE
66E	PRES.	0	VOID				NANO MARL, dusky yellow green (5GY 5/2) with interbeds of dark greenish gray (5G 4/1) and grayish olive (10Y 4/2), very stiff, slight drilling disturbance, bioturbated (Zoophycos), especially in Section 6.
	ABUND.	1.0	N				SS at 3-100 (dominant lithology)
	RADS	0.5	N				Quartz R Heavy minerals A
	FOSIL		N				Clay R Volcanic glass R
	PRES.		N				Micro nodules R
	ABUND.		N				Forams R Carbonate rhombs R
	RADS		N				Nannos R Nannos A
	FOSIL		N				Diatoms R Rad.
	PRES.		N				Sponge spicules R
	ABUND.		N				Sponge spicules R
	RADS		N				SS at 6-100 (dominant lithology)
	FOSIL		N				Quartz R Clay A
	PRES.		N				Volcanic glass R Micro nodules R
	ABUND.		N				Zeolites R Fe/Mn R
	RADS		N				Carbonate rhombs R Forams R
	FOSIL		N				Nannos R A
	PRES.		N				Diatoms R Rad.
	ABUND.		N				Sponge spicules R
	RADS		N				Fish debris R
	FOSIL		N				Carbonate Bomb: 3-114 to 115 cm = 44%
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				
	RADS		N				
	FOSIL		N				
	PRES.		N				
	ABUND.		N				</

Site 369 Hole A		Core 21 Cored Interval : 232.0-241.5 m			
ZONES	AGE	FOSILS	CHARACTER	METERS	LITHOLOGY
NANNO	PRES.	RADS	VOID	0	
NANNO	PLAT.	RADS	N	0.5	
NANNO	PLAT.	RADS	N	1.0	
NANNO	PLAT.	RADS	N	2	
NANNO	PLAT.	RADS	N	3	
DEFORMATION LITHOLOGY					
LITHO. SAMPLE					
LITHOLOGY DESCRIPTION					
NANNO MAR, dusky yellow green (SGv 5/2) with grayish blue green (BG 5/2), pale green (10G 6/2), and greenish gray (SG 6/1), very firm, slight drilling disturbance, bioturbated.					
SS at 2-100 (dominant lithology)					
Quartz R					
Volcanic glass R					
Zeolites R					
Carbonate R					
unspecified R					
Forams R					
Diatoms R					
Sponge spicules R					
SS at 5-100 (dominant lithology)					
Quartz R					
Volcanic glass R					
Micronodules R					
Carbonate R					
unspecified R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					
Radi R					
Silicoflagegulates R					
SS at 2-100 (dominant lithology)					
Quartz R					
Clay R					
Glaconite R					
Zeolites R					
Carbonate rhombs R					
Nanom R					

Site 369 Hole A		Core 22 Cored Interval: 241.5-251.0 m	
		LITHOLOGY	LITHOLOGIC DESCRIPTION
ZONES	FOSIL CHARACTER	METERS	
AEF	FORAMS	0	VOID
		0.5	N
		1.0	5G 4/1
		1	
		2	
		3	
		4	
		5	
		6	
		7	
		8	
		9	
		10	
		11	
		12	
		13	
		14	
		15	
		16	
		17	
		18	
		19	
		20	
		21	
		22	
		23	
		24	
		25	
		26	
		27	
		28	
		29	
		30	
		31	
		32	
		33	
		34	
		35	
		36	
		37	
		38	
		39	
		40	
		41	
		42	
		43	
		44	
		45	
		46	
		47	
		48	
		49	
		50	
		51	
		52	
		53	
		54	
		55	
		56	
		57	
		58	
		59	
		60	
		61	
		62	
		63	
		64	
		65	
		66	
		67	
		68	
		69	
		70	
		71	
		72	
		73	
		74	
		75	
		76	
		77	
		78	
		79	
		80	
		81	
		82	
		83	
		84	
		85	
		86	
		87	
		88	
		89	
		90	
		91	
		92	
		93	
		94	
		95	
		96	
		97	
		98	
		99	
		100	
		101	
		102	
		103	
		104	
		105	
		106	
		107	
		108	
		109	
		110	
		111	
		112	
		113	
		114	
		115	
		116	
		117	
		118	
		119	
		120	
		121	
		122	
		123	
		124	
		125	
		126	
		127	
		128	
		129	
		130	
		131	
		132	
		133	
		134	
		135	
		136	
		137	
		138	
		139	
		140	
		141	
		142	
		143	
		144	
		145	
		146	
		147	
		148	
		149	
		150	
		151	
		152	
		153	
		154	
		155	
		156	
		157	
		158	
		159	
		160	
		161	
		162	
		163	
		164	
		165	
		166	
		167	
		168	
		169	
		170	
		171	
		172	
		173	
		174	
		175	
		176	
		177	
		178	
		179	
		180	
		181	
		182	
		183	
		184	
		185	
		186	
		187	
		188	
		189	
		190	
		191	
		192	
		193	
		194	
		195	
		196	
		197	
		198	
		199	
		200	
		201	
		202	
		203	
		204	
		205	
		206	
		207	
		208	
		209	
		210	
		211	
		212	
		213	
		214	
		215	
		216	
		217	
		218	
		219	
		220	
		221	
		222	
		223	
		224	
		225	
		226	
		227	
		228	
		229	
		230	
		231	
		232	
		233	
		234	
		235	
		236	
		237	
		238	
		239	
		240	
		241	
		242	
		243	
		244	
		245	
		246	
		247	
		248	
		249	
		250	
		251	

UPPER OLIGOCENE

Porcardosphaeris distentulus Np24

Sphaerotilus optima opima

C. A. P. M.

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M

P21

Core Catcher

N A G

R C G

F C-A P-M


P21

Core Catcher

N A G

Site 369 Hole A		Core 23 Cored Interval: 251.0-260.5 m	
		LITHO. SAMPLE	DEFORATION
ZONES	FOSIL CHARACTER	METERS	LITHOLOGY
AEF	FORAMS	0	
		0.5	
		1.0	
		1	
		2	
		3	
		4	
		5	
		6	
		7	
		8	
		9	
		10	
		11	
		12	
		13	
		14	
		15	
		16	
		17	
		18	
		19	
		20	
		21	
		22	
		23	
		24	
		25	
		26	
		27	
		28	
		29	
		30	
		31	
		32	
		33	
		34	
		35	
		36	
		37	
		38	
		39	
		40	
		41	
		42	
		43	
		44	
		45	
		46	
		47	
		48	
		49	
		50	
		51	
		52	
		53	
		54	
		55	
		56	
		57	
		58	
		59	
		60	
		61	
		62	
		63	
		64	
		65	
		66	
		67	
		68	
		69	
		70	
		71	
		72	
		73	
		74	
		75	
		76	
		77	
		78	
		79	
		80	
		81	
		82	
		83	
		84	
		85	
		86	
		87	
		88	
		89	
		90	
		91	
		92	
		93	
		94	
		95	
		96	
		97	
		98	
		99	
		100	
		101	
		102	
		103	
		104	
		105	
		106	
		107	
		108	
		109	
		110	
		111	
		112	
		113	
		1	

146



Site 369 Hole A		Core 28		Core Interval: 298.5-308.0 m	
ZONES	AGE	FOSILS	CHARACTER	METERS	LITHOLOGY
FAIRAMS	LOWER OLIGOCENE	RADS	PRES.	0	0.5
MANNINS		RADS	ABUND.	1	1.0
ETRIGONIUM				2	
THEOCYRTINA				3	
GLOBULIGERINA				4	
TRICERATINA				5	
LITHOLOGIC DESCRIPTION					
NANNO MARL, dusky yellow green (5Gy 5/2) and greenish gray (5G 6/1), very firm, slight drilling disturbance, bioturbated. SS at 3-100 (dominant lithology)					
Quartz R Feldspar R Heavy minerals R Micronodules R Glaucite R Carbonate rhombs R Forams R Nannos A Diatoms R Radls R Sponge spicules R Silicoflagellates R Fish debris R					
SS at 5-100 (dominant lithology)					
Quartz R Clay R Micronodules R Zeolites R Nannos R Radls R Sponge spicules R Fish debris R					
5G 6/1					
5G 5/2					
5G 6/1					
100					
Core Catcher					

Site 369		Hole A		Core 33		Cored Interval: 346.0-355.5 m	
AGE		ZONES		FOSIL CHARACTER		LITHOLOGY	
BED		FOSILS		PRBS		MEETRES	
DEFORMATION		SECTION		PRBS		0	
SAMPLE		ABUNN.		1		0.5	
NANNO MARL, olive gray (5Y 3/2), very firm, slight drilling disturbance, bioturbated, appears to be a slump.		N		N		N	
ARGILLACEOUS NANNO LIMESTONE, light green gray (5Y 7/1) to very light gray (Nb), burrowed with various colors of brown.		N		N		1.0	
SS at 2-100 (dominant lithology)		N		N		N	
Quartz		N		N		N	
Volcanic glass		N		N		N	
Carbonate rhombs		N		N		N	
Diatoms		N		N		N	
Fish debris		N		N		N	
SS at 4-100 (dominant lithology)		N		N		N	
Clay		N		N		N	
Carbonate unspecified		N		N		N	
Forams		N		N		N	
Nannos		N		N		N	
A		N		N		N	
R		N		N		N	
A		N		N		N	
R		N		N		N	
R		N		N		N	
A		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N		N	
R		N		N			

Site 369 Hole A		Core 35		Cored interval: 365.0-374.5 m	
ZONES	FOSSIL CHARACTER	AGE	FORMS	RODS	LITHOLOGY
METE 23	SPRS.	0			ARGILLACEOUS NANNO LIMESTONE WITH MINOR PORCELLANITE AND CHERT
ABUND.	ABUND.	0.5	VOID		ARGILLACEOUS NANNO LIMESTONE, very light gray (N8) to very light greenish gray (5Gy 8/1), bioturbated, blue black (m) halos and limestone banding, color banding in limestone correlates to degree of cementation - the NO sections are considerably harder than the others. Pyrite is scattered throughout.
NANNO	-FOSSIL	1.0			pyrillanites and chert occur in the color cycles - always occurring in the NB layer.
LITHO. SAMPLE					
DEFORMATION					
LITHO. SECTION					
0					
1					
2					
3					
4					
5					
6					
7					
8					
9					
10?					
11/10?					
12/11?					
13/12?					
14/13?					
15/14?					
16/15?					
17/16?					
18/17?					
19/18?					
20/19?					
21/20?					
22/21?					
23/22?					
24/23?					
25/24?					
26/25?					
27/26?					
28/27?					
29/28?					
30/29?					
31/30?					
32/31?					
33/32?					
34/33?					
35/34?					
36/35?					
37/36?					
38/37?					
39/38?					
40/39?					
41/40?					
42/41?					
43/42?					
44/43?					
45/44?					
46/45?					
47/46?					
48/47?					
49/48?					
50/49?					
51/50?					
52/51?					
53/52?					
54/53?					
55/54?					
56/55?					
57/56?					
58/57?					
59/58?					
60/59?					
61/60?					
62/61?					
63/62?					
64/63?					
65/64?					
66/65?					
67/66?					
68/67?					
69/68?					
70/69?					
71/70?					
72/71?					
73/72?					
74/73?					
75/74?					
76/75?					
77/76?					
78/77?					
79/78?					
80/79?					
81/80?					
82/81?					
83/82?					
84/83?					
85/84?					
86/85?					
87/86?					
88/87?					
89/88?					
90/89?					
91/90?					
92/91?					
93/92?					
94/93?					
95/94?					
96/95?					
97/96?					
98/97?					
99/98?					
100/99?					
101/100?					
102/101?					
103/102?					
104/103?					
105/104?					
106/105?					
107/106?					
108/107?					
109/108?					
110/109?					
111/110?					
112/111?					
113/112?					
114/113?					
115/114?					
116/115?					
117/116?					
118/117?					
119/118?					
120/119?					
121/120?					
122/121?					
123/122?					
124/123?					
125/124?					
126/125?					
127/126?					
128/127?					
129/128?					
130/129?					
131/130?					
132/131?					
133/132?					
134/133?					
135/134?					
136/135?					
137/136?					
138/137?					
139/138?					
140/139?					
141/140?					
142/141?					
143/142?					
144/143?					
145/144?					
146/145?					
147/146?					
148/147?					
149/148?					
150/149?					
151/150?					
152/151?					
153/152?					
154/153?					
155/154?					
156/155?					
157/156?					
158/157?					
159/158?					
160/159?					
161/160?					
162/161?					
163/162?					
164/163?					
165/164?					
166/165?					
167/166?					
168/167?					
169/168?					
170/169?					
171/170?					
172/171?					
173/172?					
174/173?					
175/174?					
176/175?					
177/176?					
178/177?					
179/178?					
180/179?					
181/180?					
182/181?					
183/182?					
184/183?					
185/184?					
186/185?					
187/186?					
188/187?					
189/188?					
190/189?					
191/190?					
192/191?					
193/192?					
194/193?					
195/194?					
196/195?					
197/196?					
198/197?					
199/198?					
200/199?					
201/200?					
202/201?					
203/202?					
204/203?					
205/204?					
206/205?					
207/206?					
208/207?					
209/208?					
210/209?					
211/210?					
212/211?					
213/212?					
214/213?					
215/214?					
216/215?					
217/216?					
218/217?					
219/218?					
220/219?					
221/220?					
222/221?					
223/222?					
224/223?					
225/224?					
226/225?					
227/226?					
228/227?					
229/228?					
230/229?					
231/230?					
232/231?					
233/232?					
234/233?					
235/234?					

133

Site 369 Hole A		Core 40		Cored Interval : 412.5-422.0 m	
AGE	ZONES	FOSIL CHARACTER	METERS	SECTIION	LITHOLOGY
					LITHO-SAMPLE
					DEFORMATION
					SLUMP
					not older than Effelithus turritus effelit
					ALBIAN
					LITHOLOGIC DESCRIPTION
					ARGILLACEOUS CHALK, bluish white (SB 9/1), firm, slight drilling disturbance, bio-firme, slight "anisotropic" structures or nortles. Slump is composed of a variety of convoluted, varicolored marls, clays, and limestones.
					SS at 2-30 (dominant lithology)
					Quartz R
					Carbonate rhombs C
					unspecified R
					Nannos R
					5B 9/1
					30
					SLUMP
					Marthasterites furcatus
					CONOMICAN-SANTONIAN

Site 369 Hole A		Core 41		Cored Interval : 422.0-431.5 m	
AGE	ZONES	FOSIL CHARACTER	METERS	SECTIION	LITHOLOGY
					LITHO-SAMPLE
					DEFORMATION
					VOID
			0		
			0.5		
			1		
			1.0		
					TOP IS A SLUMP composed of clays, marls and nanno marls and chalks of various colors (from top to 2-115).
					NANNO-BEARING MARL, olive black (SY 2/1), firm, slight drilling disturbance, bio-disturbed, laminated with dark greenish gray (56 4/1).
					Pyrite stringer occurs at 4-52 cm.
					Barite and dolomite (?) crystals occur at 4-137 to 148.
					SLUMP
					SS at 3-40 (dominant lithology)
					Quartz R
					Feldspar A
					Clay A
					Heavy minerals R
					Carbonate R
					Forams R
					unspecified C
					Nannos C
					Core Catcher

Site 369		Hole A	Core 44	Cored Interval : 460.5-460.0 m
ZONES	FORMS	FOSIL CHARACTER	METER	LITHOLOGY
Parahabdoidithus angustatus UPPER ALBIAN				
AGE				
NANNO				
RADS				
FOSSIL				
ABUND.				
PRES.				
SECTIION				
METERS				
DEFORMATION				
LITHO. SAMPLE				
Parahabdoidithus angustatus UPPER ALBIAN-LOWER ALBIAN				
AGE				
NANNO				
RADS				
FOSSIL				
ABUND.				
PRES.				
SECTIION				
METERS				
DEFORMATION				
LITHO. SAMPLE				
Parahabdoidithus angustatus UPPER ALBIAN				
AGE				
NANNO				
RADS				
FOSSIL				
ABUND.				
PRES.				
SECTIION				
METERS				
DEFORMATION				
LITHO. SAMPLE				

Site 369		Hole A	Core 45	Cored Interval : 460.0-469.5 m
ZONES	FORMS	FOSIL CHARACTER	METER	LITHOLOGY
Parahabdoidithus angustatus UPPER ALBIAN				
AGE				
NANNO				
RADS				
FOSSIL				
ABUND.				
PRES.				
SECTIION				
METERS				
DEFORMATION				
LITHO. SAMPLE				
Parahabdoidithus angustatus UPPER ALBIAN				
AGE				
NANNO				
RADS				
FOSSIL				
ABUND.				
PRES.				
SECTIION				
METERS				
DEFORMATION				
LITHO. SAMPLE				

DEEP SEA DRILLING PROJECT

LEG 41 SITE 370

SITE SUMMARY SHEET

PRINCIPAL RESULTS:

Deep Basin off Morocco

Nearly 1200 meters of the sediment section of the oceanic basin off the Moroccan continental margin was sampled. The uppermost section, down to 420 meters, was very sparsely cored but below that level coring was closely spaced until time constraints forced us to abandon the site. The sediments are predominantly hemipelagic containing silt and sand layers and occasionally conglomerates; most of these layers probably are turbidites. The section reflects an evolution different from that of the Cape Verde Basin and of the North American Basin because here a strong terrigenous influence was felt from the lower Cretaceous to the Pleistocene. The section can be summarized as follows (from top to bottom): 1) Pleistocene clay with nannofossil and foraminifera; 2) Pleistocene to middle Miocene clay and silty nannofossil marls interbedded with silt and sand; 3) middle Miocene to late Paleocene clays with occasional radiolarians, chert, silt, sand and gravel; 4) late to early Paleocene nannofossil marls and clays; 5) early Cenomanian to Berriasian calcareous silty claystones interbedded with siltstones, sandstone and conglomerates. A hiatus of 35 m.y. separates late Cenomanian from early Paleocene sediments. No well-defined black shales were found in Aptian to Cenomanian sediments although the claystones are dark in this interval and have organic carbon contents reaching 5%. Rosettes of barite partially replaced by calcite, as at Site 369, might indicate migration of calcium sulfate-rich solutions originating from possible underlying evaporites. Calcite cementation of sandstone and siltstone beds below about 835 meters might be also related to such migration.

Date occupied	0100 March 29, 1975
Date departed	0110 April 8, 1975
Time on site	10 days, and 10 minutes
Position: Latitude	32°50.2'N
Longitude	10°46.6'W
Water depth (sea level)	4214 corrected meters
Water depth (rig floor)	4224 corrected meters
Penetration	1176.5 meters

LEG 41 SITE 370
SITE SUMMARY SHEET, con't.

Number of holes	1
Number of cores	51
Total length of cored section	483 meters
Total core recovered	202.7 meters
Percentage core recovery	42%

Oldest Sediment Cored

Depth subbottom	1176.5 meters
Nature	Claystone and siltstone
Age	Berriasian
Measured velocity	3.82 km/sec

Site 370 Hole Core 5 Cored Interval: 321.5-331.0 m									
LITHOLOGIC DESCRIPTION									
AGE	FORMS	NANOS	RADS	SECT. ZONES	CHARACTER	FOSIL	METERS	PRBS.	LITHOLOGY
				0	INTERBEDDED CALCAREOUS SILTY CLAY, SILTY CLAY, AND SANDY NANNO MARL WITH OCCASIONAL TURBIDITES.				
				1	CALCAREOUS SILTY CLAY, olive gray (SY 4/1), firm, slight drilling disturbance, bioturbated, laminated.		0.5	N	SILTY CLAY, olive gray (SY 4/1), firm, slight drilling disturbance, laminated with greenish gray (SY 6/1).
				2	SILTY CLAY, greenish gray (SY 6/1), firm, slight drilling disturbance, thin dark gray laminae. Occurs as basal and part of above unit 5-40 cm thick.		1.0	N	SILTY CLAY, dark greenish gray (SY 4/1), firm to semi-firified, slight drilling disturbance, parallel and irregular faint laminations.
				3	Indicates sand bed.				PORCELLANITE, dark gray (N3), occurs in the upper part of the core.
				4	SANDY NANNO MARL, greenish gray (SY 6/1), massive, basal A units of TURBIDITE.				SS at 1-63 (dominant lithology)
				5	SY 4/1				Quartz
				6	SY 6/1				Feldspar
				7	60				Glaucite
				8	SY 4/1				Rhomb
				9	SY 6/1				unspecified
				10	SS at 1-10 (dominant lithology)				C
				11	Quartz				Rads
				12	SY 4/1				Fish debris
				13	SY 6/1				SS at 2-73 (dominant lithology)
				14	Quartz				Clay
				15	SY 4/1				Carbonate
				16	SY 6/1				Rhomb
				17	SS at 2-60 (dominant lithology)				unspecified
				18	Quartz				R
				19	SY 6/1				Nanos
				20	INTERBEDDED				Rhomb
				21	SY 4/1				unspecified
				22	SS at 4-50 (dominant lithology)				Nanos
				23	Quartz				A
				24	SY 4/1				Glaucite
				25	SY 6/1				Carbonate
				26	INTERBEDDED				Rhomb
				27	SY 4/1				unspecified
				28	SS at 4-50 (dominant lithology)				Rhomb
				29	Quartz				unspecified
				30	SY 6/1				Rhomb
				31	INTERBEDDED				unspecified
				32	SY 4/1				Rhomb
				33	SS at 4-50 (dominant lithology)				unspecified
				34	Quartz				Rhomb
				35	SY 6/1				unspecified
				36	INTERBEDDED				Rhomb
				37	SY 4/1				unspecified
				38	SS at 4-50 (dominant lithology)				Rhomb
				39	Quartz				unspecified
				40	SY 6/1				Rhomb
				41	INTERBEDDED				unspecified
				42	SY 4/1				Rhomb
				43	SS at 4-50 (dominant lithology)				unspecified
				44	Quartz				Rhomb
				45	SY 6/1				unspecified
				46	INTERBEDDED				Rhomb
				47	SY 4/1				unspecified
				48	SS at 4-50 (dominant lithology)				Rhomb
				49	Quartz				unspecified
				50	SY 6/1				Rhomb
				51	INTERBEDDED				unspecified
				52	SY 4/1				Rhomb
				53	SS at 4-50 (dominant lithology)				unspecified
				54	Quartz				Rhomb
				55	SY 6/1				unspecified
				56	INTERBEDDED				Rhomb
				57	SY 4/1				unspecified
				58	SS at 4-50 (dominant lithology)				Rhomb
				59	Quartz				unspecified
				60	SY 6/1				Rhomb
				61	INTERBEDDED				unspecified
				62	SY 4/1				Rhomb
				63	SS at 4-50 (dominant lithology)				unspecified
				64	Quartz				Rhomb
				65	SY 6/1				unspecified
				66	INTERBEDDED				Rhomb
				67	SY 4/1				unspecified
				68	SS at 4-50 (dominant lithology)				Rhomb
				69	Quartz				unspecified
				70	SY 6/1				Rhomb
				71	INTERBEDDED				unspecified
				72	SY 4/1				Rhomb
				73	SS at 4-50 (dominant lithology)				unspecified
				74	Quartz				Rhomb
				75	SY 6/1				unspecified
				76	INTERBEDDED				Rhomb
				77	SY 4/1				unspecified
				78	SS at 4-50 (dominant lithology)				Rhomb
				79	Quartz				unspecified
				80	SY 6/1				Rhomb
				81	INTERBEDDED				unspecified
				82	SY 4/1				Rhomb
				83	SS at 4-50 (dominant lithology)				unspecified
				84	Quartz				Rhomb
				85	SY 6/1				unspecified
				86	INTERBEDDED				Rhomb
				87	SY 4/1				unspecified
				88	SS at 4-50 (dominant lithology)				Rhomb
				89	Quartz				unspecified
				90	SY 6/1				Rhomb
				91	INTERBEDDED				unspecified
				92	SY 4/1				Rhomb
				93	SS at 4-50 (dominant lithology)				unspecified
				94	Quartz				Rhomb
				95	SY 6/1				unspecified
				96	INTERBEDDED				Rhomb
				97	SY 4/1				unspecified
				98	SS at 4-50 (dominant lithology)				Rhomb
				99	Quartz				unspecified
				100	SY 6/1				Rhomb
				101	INTERBEDDED				unspecified
				102	SY 4/1				Rhomb
				103	SS at 4-50 (dominant lithology)				unspecified
				104	Quartz				Rhomb
				105	SY 6/1				unspecified
				106	INTERBEDDED				Rhomb
				107	SY 4/1				unspecified
				108	SS at 4-50 (dominant lithology)				Rhomb
				109	Quartz				unspecified
				110	SY 6/1				Rhomb
				111	INTERBEDDED				unspecified
				112	SY 4/1				Rhomb
				113	SS at 4-50 (dominant lithology)				unspecified
				114	Quartz				Rhomb
				115	SY 6/1				unspecified
				116	INTERBEDDED				Rhomb
				117	SY 4/1				unspecified
				118	SS at 4-50 (dominant lithology)				Rhomb
				119	Quartz				unspecified
				120	SY 6/1				Rhomb
				121	INTERBEDDED				unspecified
				122	SY 4/1				Rhomb
				123	SS at 4-50 (dominant lithology)				unspecified
				124	Quartz				Rhomb
				125	SY 6/1				unspecified
				126	INTERBEDDED				Rhomb
				127	SY 4/1				unspecified
				128	SS at 4-50 (dominant lithology)				Rhomb
				129	Quartz				unspecified
				130	SY 6/1				Rhomb
				131	INTERBEDDED				unspecified
				132	SY 4/1				Rhomb
				133	SS at 4-50 (dominant lithology)				unspecified
				134	Quartz				Rhomb
				135	SY 6/1				unspecified
				136	INTERBEDDED				Rhomb
				137	SY 4/1				unspecified
				138	SS at 4-50 (dominant lithology)				Rhomb
				139	Quartz				unspecified
				140	SY 6/1				Rhomb
				141	INTERBEDDED				unspecified
				142	SY 4/1				Rhomb
				143	SS at 4-50 (dominant lithology)				unspecified
				144	Quartz				Rhomb
				145	SY 6/1				unspecified
				146	INTERBEDDED				Rhomb
				147	SY 4/1				unspecified
				148	SS at 4-50 (dominant lithology)				Rhomb
				149	Quartz				unspecified
				150	SY 6/1				Rhomb
				151	INTERBEDDED				unspecified
				152	SY 4/1				Rhomb
				153	SS at 4-50 (dominant lithology)				unspecified
				154	Quartz				Rhomb
				155	SY 6/1				unspecified
				156	INTERBEDDED				Rhomb
				157	SY 4/1				unspecified
				158	SS at 4-50 (dominant lithology)				Rhomb
				159	Quartz				unspecified
				160	SY 6/1				Rhomb
				161	INTERBEDDED				unspecified
				162	SY 4/1				Rhomb
				163	SS at 4-50 (dominant lithology)				unspecified
				164	Quartz				Rhomb
				165	SY 6/1				unspecified
				166	INTERBEDDED				Rhomb
				167	SY 4/1				unspecified
				168	SS at 4-50 (dominant lithology)				Rhomb
				169	Quartz				unspecified
				170	SY 6/1				Rhomb
				171	INTERBEDDED				unspecified
				172	SY 4/1				Rhomb
				173	SS at 4-50 (dominant lithology)				unspecified
				174	Quartz				Rhomb
				175	SY 6/1				unspecified
				176	INTERBEDDED				Rhomb
				177	SY 4/1				unspecified
				178	SS at 4-50 (dominant lithology)				Rhomb
				179	Quartz				unspecified
				180	SY 6/1				Rhomb
				181	INTERBEDDED				unspecified
				182	SY 4/1				Rhomb
				183	SS at 4-50 (dominant lithology)				unspecified
				184	Quartz				

Site 370		Hole	Core 8	Cored Interval: 464.0-473.5 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION
MIDDLE EOCENE	CHTHIAGMOLITHUS ALATUS - DISCOASTER TANINODIFER NP15-NP16			SILTY CLAY OVERLYING CLAYSTONE WITH MINOR PORCELLANITE THROUGHOUT.
				PORCELLANITES, olive gray (SY 4/1).
				SECTION 1: SILTY CLAY, dark greenish gray (SY 4/1), firm to semilithified, no drilling disturbance, parallel and irregular faint laminations.
				SECTION 2: CLAYSTONE, olive gray (SY 3/2), lithified, no drilling disturbance, massive, bioturbated streaks and laminae of pale green (10G 6/2). MICROCONGLOMERATE, at Section 2-14 cm consists of grains of brownish limestone (up to 2 mm). Basalt contact is sharp with scouring.
				MANGNO-BEADING CLAYSTONE at Section 2-12 to 115 cm, yellowish gray (SY 8/1), lithified, bioturbated, sharp boundaries.
				SS at 1-125 (dominant lithology) Quartz R Glaucite R Carbonate R Nannos R Radls R Unspecified R Forams R Nannos C
				SS at 2-114 (minor lithology) Quartz R Clay R Carbonate D Unspecified R Nannos R
				Carbone Bomb: 2-82 to 83 cm = 19%

Site 370		Hole	Core 9	Cored Interval: 483.0-492.5 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION
MIDDLE EOCENE	CHTHIAGMOLITHUS ALATUS - DISCOASTER TANINODIFER NP15-NP16			SILTY CLAY OVERLYING CLAYSTONE WITH MINOR PORCELLANITE THROUGHOUT.
				PORCELLANITES, olive gray (SY 4/1).
				SECTION 1: CLAYSTONE, grayish olive green (SY 4/1), lithified, tanned, very rare disturbance, a chevron fold occurs at ~30 cm.
				CLAYSTONE WITH MINOR PORCELLANITE, PORCELLANITES, olive gray (SY 4/1).
				SS at 93 Quartz R Feldspar R Glaucite R Micronodules R Carbonate rhombs R Unspecified R Fish debris R
				SS at CC Quartz R Glaucite R Micronodules R Carbonate rhombs R Unspecified R Fish debris R
				Carbonate Bomb: 1-122 to 123 cm = 13%

Site 370		Hole	Core 10	Cored Interval: 502.0-511.5 m
AGE	ZONES	FOSIL CHARACTER	LITHOLOGY	LITHOLOGIC DESCRIPTION
MIDDLE EOCENE	CHTHIAGMOLITHUS ALATUS - DISCOASTER TANINODIFER NP15-NP16			SILTY CLAYSTONE WITH MINOR PORCELLANITE.
				PORCELLANITES, at top of core, olive gray (SY 4/1), encloses some surrounded pebbles of brown mudstone (up to 2 cm in size).
				SS at 2-20 (dominant lithology) Quartz C Feldspar R Micronodules R Carbonate rhombs R Unspecified R Fish debris R
				SS at CC Quartz R Glaucite R Micronodules R Carbonate rhombs R Unspecified R Fish debris R
				Carbonate Bomb: 2-50 to 51 cm = 15%

Site 370 Hole		Core 11		Cored Interval: 521.0-530.5 m		Site 370 Hole		Core 13		Cored Interval: 549.5-559.0 m	
AGES	ZONES	FOSIL CHARACTER	LITHOLOGY	METERS	SECTION	AGES	ZONES	FOSIL CHARACTER	LITHOLOGY	METERS	SECTION
LITHOLOGIC DESCRIPTION											
			1 cm of CALCAEUS SILTY CLAYSTONE, light olive gray (5Y 5/2).	0							
LITHOLOGIC DESCRIPTION											
			CALCAREOUS SILTY CALSTONE WITH MINOR LIMESTONE AND PORCELLANITE, CALCAREOUS SILY CLAYSTONE, grayish olive (5Y 4/2), lithified, moderately bio-turbated, scattered sand-sized porcellanite, irregular laminae of fine sand grains composed of mudstone cemented by calcite, rare clay laminae, LIMESTONE, microcrystalline, bluish white (SB 9/1).								
			PORCELLANITE, olive gray (5Y 4/1).								
LITHOLOGIC DESCRIPTION											
			SS at 2-50 (dominant lithology)								
			Quartz C Feldspar R								
			Heavy minerals A								
			Clay R								
			Glaucite R								
			Fish debris R								
			Micro nodules R								
			Carbonate rhombs R								
			Forams R								
			Unspecified C								
			Nannos R								
			Sponge spicules R								
LITHOLOGIC DESCRIPTION											
			SS at 2-56 (minor lithology)								
			Quartz C Feldspar R								
			Heavy minerals R								
			Clay R								
			Glaucite R								
			Fish debris R								
			Micro nodules R								
			Carbonate rhombs R								
			Forams R								
			Unspecified C								
			Nannos R								
			Sponge spicules R								
LITHOLOGIC DESCRIPTION											
			SS at 2-97 (minor lithology)								
			Clay C Carbonate rhombs R								
			Unspecified D								
			Nannos R								
LITHOLOGIC DESCRIPTION											
			SS at CC								
			Quartz R								
			Feldspar R								
			Clay A								
			Heavy minerals R								
			Pyrite R								
			Glaucite R								
			Micro nodules R								
			Zeolites R								
			Carbonate rhombs C								
			Forams R								
			Unspecified R								
			Nannos R								
			Fish debris R								
LITHOLOGIC DESCRIPTION											
			Carbone Bomb 2-10 to 102 cm = 70%								
			2-75 to 76 cm = 9%								
LITHOLOGIC DESCRIPTION											
			PEBBLE CONGLOMERATE, olive gray (5Y 4/1), pebbles rounded-subangular, size 1 to 2 cm, composed of porcellanite, mudstone, glauconite, and rare forams, all cemented with silica.								
			CONGLOMERATE consists of pebbles and cement within the conglomerate and of scattered rounded pebbles near the base of Section 2.								
LITHOLOGIC DESCRIPTION											
			SS at 2-10 (minor lithology)								
			Quartz C Feldspar R								
			Heavy minerals R								
			Clay A								
			Glaucite R								
			Pyrite R								
			Carbonate R								
			Unspecified C								
			Fish debris R								
			Mollusc debris R								
LITHOLOGIC DESCRIPTION											
			SS at 2-100 (dominant lithology)								
			Quartz R Feldspar R								
			Heavy minerals R								
			Clay A								
			Glaucite R								
			Pyrite R								
			Carbonate R								
			Unspecified C								
			Fish debris R								
			Mollusc debris R								
LITHOLOGIC DESCRIPTION											
			SS at 2-100 (dominant lithology)								
			Quartz R Feldspar R								
			Heavy minerals R								
			Clay A								
			Glaucite R								
			Pyrite R								
			Carbonate R								
			Unspecified C								
			Fish debris R								
			Mollusc debris R								

1461

Site 370 Hole		Core 14 Cored Interval : 568.5-578.0 m		Site 370 Hole		Core 15 Cored Interval : 587.5-597.0 m	
LITHOLOGIC DESCRIPTION		LITHOLOGIC DESCRIPTION		LITHOLOGIC DESCRIPTION		LITHOLOGIC DESCRIPTION	
AGE	FORMS	ROADS	NAME	AGE	FORMS	ROADS	NAME
ZONES	FOSSTIL	PRDS.	SECTON	ZONES	FOSSTIL	PRDS.	SECTON
METERS	CHARACTER	ABUND.		METERS	CHARACTER	ABUND.	
0	VOID	0.5		0	VOID	0.5	
CALCAREOUS SILTY CLAYSTONE WITH MINOR CHERT; OVERLYING CALCAREOUS CLAYEY SILTSTONE TO SANDSTONE.		CALCAREOUS SILTY CLAYSTONE, olive gray (SY 3/2), moderately bioturbated with occasional Zoothipcos, thin silt laminae siliceous at 2-25 to 40, 2-60 to 63, 2-75 to 123.		CALCAREOUS CLAYEY SILTSTONE AND SANDSTONE below 4.80, dark yellowish brown (Tyr 4/2), consists of brownish grains of claystone with rare forams, cemented by calcite.		CALCAREOUS CLAYEY SILTSTONE AND SANDSTONE below 4.80, dark yellowish brown (Tyr 4/2), consists of brownish grains of claystone with rare forams, cemented by calcite.	
0.5		0.5		0.5		0.5	
1		1		1		1	
1.0		1.0		1.0		1.0	
2		2		2		2	
1.5		1.5		1.5		1.5	
3		3		3		3	
3.0		3.0		3.0		3.0	
4		4		4		4	
4.0		4.0		4.0		4.0	
5		5		5		5	
5.0		5.0		5.0		5.0	
5.5		5.5		5.5		5.5	
6		6		6		6	
6.5		6.5		6.5		6.5	
7		7		7		7	
7.5		7.5		7.5		7.5	
8		8		8		8	
8.5		8.5		8.5		8.5	
9		9		9		9	
9.5		9.5		9.5		9.5	
10		10		10		10	
10.5		10.5		10.5		10.5	
11		11		11		11	
11.5		11.5		11.5		11.5	
12		12		12		12	
12.5		12.5		12.5		12.5	
13		13		13		13	
13.5		13.5		13.5		13.5	
14		14		14		14	
14.5		14.5		14.5		14.5	
15		15		15		15	
15.5		15.5		15.5		15.5	
16		16		16		16	
16.5		16.5		16.5		16.5	
17		17		17		17	
17.5		17.5		17.5		17.5	
18		18		18		18	
18.5		18.5		18.5		18.5	
19		19		19		19	
19.5		19.5		19.5		19.5	
20		20		20		20	
20.5		20.5		20.5		20.5	
21		21		21		21	
21.5		21.5		21.5		21.5	
22		22		22		22	
22.5		22.5		22.5		22.5	
23		23		23		23	
23.5		23.5		23.5		23.5	
24		24		24		24	
24.5		24.5		24.5		24.5	
25		25		25		25	
25.5		25.5		25.5		25.5	
26		26		26		26	
26.5		26.5		26.5		26.5	
27		27		27		27	
27.5		27.5		27.5		27.5	
28		28		28		28	
28.5		28.5		28.5		28.5	
29		29		29		29	
29.5		29.5		29.5		29.5	
30		30		30		30	
30.5		30.5		30.5		30.5	
31		31		31		31	
31.5		31.5		31.5		31.5	
32		32		32		32	
32.5		32.5		32.5		32.5	
33		33		33		33	
33.5		33.5		33.5		33.5	
34		34		34		34	
34.5		34.5		34.5		34.5	
35		35		35		35	
35.5		35.5		35.5		35.5	
36		36		36		36	
36.5		36.5		36.5		36.5	
37		37		37		37	
37.5		37.5		37.5		37.5	
38		38		38		38	
38.5		38.5		38.5		38.5	
39		39		39		39	
39.5		39.5		39.5		39.5	
40		40		40		40	
40.5		40.5		40.5		40.5	
41		41		41		41	
41.5		41.5		41.5		41.5	
42		42		42		42	
42.5		42.5		42.5		42.5	
43		43		43		43	
43.5		43.5		43.5		43.5	
44		44		44		44	
44.5		44.5		44.5		44.5	
45		45		45		45	
45.5		45.5		45.5		45.5	
46		46		46		46	
46.5		46.5		46.5		46.5	
47		47		47		47	
47.5		47.5		47.5		47.5	
48		48		48		48	
48.5		48.5		48.5		48.5	
49		49		49		49	
49.5		49.5		49.5		49.5	
50		50		50		50	
50.5		50.5		50.5		50.5	
51		51		51		51	
51.5		51.5</td					

Site 370 Hole		Core 17		Cored Interval: 616.0-625.5 m	
				LITHOLOGIC DESCRIPTION	
ZONES	FOSIL CHARACTER	SECTION	METERS	LITHOLOGY	
ABUND.	RADS	PRES.			
FOAMS	NANOS				
RODS					
LOWER EOCENE		AGE		LITHO. SAMPLE	
Globorotalia aragonensis - Globorotalia palmerae F8-P9		Marthesasterites tridracchatus - Discocaster subtidianensis NP12-NP14		CALCAREOUS NANO-BEARING CLAYSTONE, INTERBEDDED WITH SILTY CLAYSTONE AND CLAYEY SILTSTONE/SANDSTONE, WITH CHERT.	
				CALCAREOUS NANO-BEARING CLAYSTONE, light olive gray (5Y 7/2) with pale blue green (5BG 7/2) calcareous silty claystone laminae (0.5 to 1 cm thick), slightly bioturbated. SILTY CLAY AND CLAYEY SILT laminae occur throughout.	
		CHERT, occurs as nodules and interbeds.			
				SS at 1-147 (dominant lithology)	
		Quartz		Quartz	
		Clay		Pyrite	
		Carbonate rhombs		Carbonate rhombs	
		Nanosc.		Forams	
		unspecified		Nannos	
				Rads	
				Chaledony	
				Sponge spicules	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay		Heavy minerals	
		Carbonate		Glaucocrite	
		unspecified		Micromoldes	
				Carbonate rhombs	
				Nannos	
				Fish debris	
				unspecified	
				C	
				SS at 2-16 (dominant lithology)	
		Quartz		Quartz	
		Clay</			

UPPER ALBIAN TO LOWER CENOMANTAN		ROTALIPORA APENNINICA		EFFELLIUS TURFSEEFFELI		Core N	
AGE	ZONES	FOSSSL	RADS	MAMS	EFFL	EFFL	CATCHER
Site 370	Hole	0	0.5	0.5	N	N	N
Core 22	LITHOLOGY	VOID					
Cored Interval : 692.0-701.5 m	METERS	0	1.0	1.0			
	SECTION	0	1	1			
	PRES.						
	ASUND.						
	FOSSSL						
	RADS						
	MAMS						
	DEFORMATION						
	LITHO. SAMPLE						
	LITHOLOGY						
	METERS						
	0	0.5	1.0	1.0	2	3	4
	1						
	2						
	3						
	4						
	48						

Site 370	Hole	Core 20	Cored Interval: 673.0-682.5 m	LITHOLOGIC DESCRIPTION										LITHO. SAMPLE		
				AGE	FERRAMS			RADAS			NANNO			FERRAMS		
ZONES	CHARACTER	ABUND.	FOSSL	SECTN	PRBS.	PRBS.	SECTN	PRBS.	PRBS.	SECTN	PRBS.	PRBS.	SECTN	PRBS.	DEFORMATION	LITHO. SAMPLE
				0				0.5		0			0		VOID	NANNO BEARING CLAYSTONE AND AG ILLACCEOUS LIMESTONE.
				1	N			1.0		1	N		1			NANNO-BEARING CLAYSTONE, dark greenish gray (5G 4/1), firm, slight drilling disturbance, homogeneous, pyrite rare.
				2	N					2	N		2			MICROCRYSTALLINE ARGILLACEOUS LIMESTONE, light greenish gray (5G 6/1) to light gray (NB), wavy laminae, gypsum or barite rosettes and laminae at 2-103 to 110 cm.
				3	N					3	N		3			SS at 1-126 (dominant lithology)
																Quartz R
																Feldspar R
																Clay A
																Volcanic glass R
																Fish debris R
																Pyrite R
																Micronodular R
																Carbonate rhombs R
																Forams C
																Nannos R
																SS at 2-115 (minor lithology)
																Clay C
																Carbonate D
																unspecified
																Carbo. Bomb: 2-72 to 73 cm = 11%

148

UPPER APPITAN TO LOWER ALBIAN										PARAHEDDOLITES ANGSTUTUS											
Core 31		Cored Interval : 825.0-824.5 m		Hole Site 370		Fossil Character		Fossils		Rocks		Section		Metres		Lithology		Litho. Sample		Description	
AEE		Forms		Morphos		Fossils		Pres.		Rocks		Section		Metres		Lithology		Litho. Sample		DEFORMATION	
0		N		N		N		R		N		N		0.5		56Y 4/1		SS at 2/1		Alternating Manganese-bearing Claystone and Silty Claystone (5Y 2/1) with varying silt content, in places bioturbated, some intervals thinly laminated with silt, and sand occasionally graded. A slump occurs at 1-75 to 100 cm.	
															68		56Y 4/1		SS at 2/1		Silty Claystone, dark greenish gray (56Y 4/1), bioturbated with thin horrows (fucoids).
															87		56Y 4/1		SS at 1-68 (dominant lithology)		Quartz C Feldspar
															5Y 2/1		Heavy minerals R		Quartz C Feldspar		Pyrite R Clay
															5Y 2/1		Pyrite R		Pyrite R Clay		Carbonate rhombs R
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		Unspecified R
															5Y 2/1		Fish debris R		Fish debris R		
															52		SS at 1-81 (minor lithology)		SS at 1-81 (minor lithology)		
															140		Quartz C Feldspar		Quartz C Feldspar		Pyrite R Clay
															5Y 2/1		Glaucocrite R		Glaucocrite R		Pyrite R Clay
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		Unspecified R
															5Y 2/1		Nannos R		Nannos R		Fish debris R
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 1-140 (minor lithology)		SS at 1-140 (minor lithology)		
															5Y 2/1		Quartz C		Quartz C		
															5Y 2/1		Pyrite R		Pyrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Nannos C		Nannos C		Unspecified R
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 2-52 (minor lithology)		SS at 2-52 (minor lithology)		
															5Y 2/1		Quartz C		Quartz C		
															5Y 2/1		Pyrite R		Pyrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos R		Nannos R		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-42 (dominant lithology)		SS at 4-42 (dominant lithology)		
															5Y 2/1		Quartz C		Quartz C		
															5Y 2/1		Pyrite D		Pyrite D		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		
															5Y 2/1		Glaucocrite R		Glaucocrite R		
															5Y 2/1		Carbonate rhombs R		Carbonate rhombs R		
															5Y 2/1		Unspecified R		Unspecified R		
															5Y 2/1		Nannos C		Nannos C		
															5Y 2/1		Plant debris R		Plant debris R		
															5Y 2/1		SS at 4-59 (minor lithology)		SS at 4-59 (minor lithology)		
															5Y 2/1		Quartz A		Quartz A		

Site 370		Hole	Core 41	Cored Interval : 986.5-986.0 m
				LITHOLOGIC DESCRIPTION
AGE	ZONES	FOSIL CHARACTER	METRES	LITHOLOGY
			0	NANNO MARLSTONE INTERBEDDED WITH THIN LAYERS OF ARGILLACEOUS CALCAREOUS SILSTONE/SANDSTONE.
				NANNO MARLSTONE, grayish green (SGY 5/2), interbedded with grayish olive (SGY 3/2).
				Thin laminae of pale yellowish brown (TGR 6/2) ARGILLACEOUS SILTSTONE and SANDSTONE, rare bioturbation.
			0.5	SS at 1-37 (dominant lithology)
				Quartz R Clay A
				Pyrite R Carbonate R
				Carbonate rhombs R Unspecified R
				Nannos A
			1	SS at 1-54 (dominant lithology)
				Quartz R Feldspar R
				Clay A
				Carbonate rhombs R
				Unspecified R
				Pyrite R
				Plant debris R
			1.0	SS at 2-64 (minor lithology)
				Quartz C Feldspar R
				Heavy minerals R
				Clay A
				Glaucite R
				Carbonate R
				Unspecified A
			2	SS at 4-27 (minor lithology)
				Quartz C Feldspar R
				Heavy minerals R
				Clay R
				Glaucite R
				Carbonate R
				Unspecified A
			3	SS at 4-39 (minor lithology)
				Quartz C Feldspar R
				Heavy minerals R
				Clay R
				Glaucite R
				Carbonate R
				Unspecified A
			4	R - Core Catcher
				R - N C W F R P
			27	

Site 370		Hole	Core 42	Cored Interval : 1005.5-1015.0 m
				LITHOLOGIC DESCRIPTION
AGE	ZONES	FOSIL CHARACTER	METRES	LITHOLOGY
			0	NANNO-BEARING MARLSTONE INTERBEDDED WITH CALCAREOUS SILSTONE/SANDSTONE AND CLAYSTONE.
				NANNO-BEARING MARLSTONE, olive gray (SGY 4/1), with a few thin CALCAREOUS SILSTONE AND SANDSTONE which are laminated and cross-bedded, rare convolute laminations.
				CLAYSTONE, pale brown (SGY 5/2), organic matter scattered throughout parallel to bedding planes.
				Claystone occurs at: 2-65 to 70, 2-120 to 126, 3-23 to 25, 3-36 to 42, 3-74 to 76, 3-126 to 130, 4-38 to 45, 4-71 to 76, 4-90 to 100.
			1	VOID
			1.0	
			2	SS at 2-74 (dominant lithology)
				Quartz R Heavy minerals R
				Clay A
				Carbonate rhombs R
				Unspecified R
				Plant debris C
			3	SS at 4-38 (minor lithology)
				Quartz R Heavy minerals R
				Clay D
				Carbonate rhombs R
				Unspecified R
				Fish debris R
			4	SS at 4-80 (dominant lithology)
				Quartz R Heavy minerals R
				Clay R
				Pyrite R
				Carbonate rhombs R
				Nannos R
				Unspecified R
				Plant debris R
			5	SS at 4-93 (minor lithology)
				Quartz C Feldspar R
				Heavy minerals R
				Clay A
				Glaucite R
				Pyrite R
				Nannos R
				Unspecified R
				Fish debris R

Site 370 Hole Core 44		Cored Interval: 1043.5-1053.0 m	
AGE	ZONES	FOSIL CHARACTER	SECTORS
R	FORAMS RHAMNOS	ABUND. FREQU.	PRES.
N	FORAMS RHAMNOS	ABUND. FREQU.	PRES.
F	FORAMS RHAMNOS	ABUND. FREQU.	PRES.
M	FORAMS RHAMNOS	ABUND. FREQU.	PRES.
G	FORAMS RHAMNOS	ABUND. FREQU.	PRES.
LITHOLOGY			
0		VOID	
1	0.5		38
1.0			
2			
3			
DEFORMATION			
LITHO. SAMPLE			
LITHOLOGIC DESCRIPTION			
SILTY MUD-BEARING CLAYSTONE WITH INTERBEDDED CALCIAREOUS ARGILLACEOUS SILTSTONE.			
SILTY MUD-BEARING CLAYSTONE (SS 5/2) to olive gray (SS 4/7), laminated.			
CALCIAREOUS ARGILLACEOUS SILTSTONE, medium gray (NS).			
SS at 1-38 (dominant lithology)			
Quartz C Heavy minerals R			
Clay A Pyrite R			
Carbonate Rhombs R			
unspecified Nannos R			
Fish debris R			
SS at 1-50 (dominant lithology)			
Quartz C Feldspar R			
Heavy minerals Clay A			
Glaconite Pyrite R			
Carbonate Rhombs R			
unspecified Nannos C			
SS at 1-107 (minor lithology)			
Quartz A Clay A			
Glaconite R Pyrite R			
Carbonate Rhombs R			
unspecified R Nannos R			
Fish debris R			
SS at 3-31 (minor lithology)			
Quartz C Feldspar R			
Heavy minerals Clay A			
Glaconite Pyrite R			
Carbonate Rhombs R			
unspecified Nannos R			
Fish debris R			
Core Catcher			
NEOCOMIAN			

Site 370		Core 45		Cored Interval : 1062.5-1072.0 m	
Hole	ZONES	FOSIL CHARACTER	LITHOLOGY	METRES	LITHO. SAMPLE
			VOID	0	
				0.5	CLAYSTONE WITH INTERBEDDED SILSTONE.
				1	CLAYSTONE, grayish green (56 5/2) and grayish brown (5YR 3/2), with inter- laminations of olive gray (5Y 4/1) variable nanno content.
				1.0	CALCAREOUS ARGILLACEOUS SILSTONES, laminated.
				2	SS at 2-30 (dominant lithology)
				30	Quartz R Clay D Carbonate R unspecified R Fish debris R
				31	SS at 3-131 (dominant lithology)
				32	Quartz R Clay A Fish debris R Carbonate R unspecified R
				33	SS at 4-53 (dominant lithology)
				34	Quartz R Clay D Carbonate R unspecified R Fish debris R
				53	NEOCOMIAN
				131	Core Catcher
					R F C P F

